



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 401/06, 401/14, 211/22, 205/04, 403/04, A61K 31/445	A1	(11) International Publication Number: WO 99/01451 (43) International Publication Date: 14 January 1999 (14.01.99)
(21) International Application Number: PCT/EP98/04177 (22) International Filing Date: 1 July 1998 (01.07.98) (30) Priority Data: 9714129.5 4 July 1997 (04.07.97) GB (71) Applicant (for GB only): PFIZER LIMITED [GB/GB]; Ramsgate Road, Sandwich, Kent CT13 9NJ (GB). (71) Applicant (for all designated States except GB US): PFIZER INC. [US/US]; 235 East 42nd Street, New York, NY 10017 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): ALKER, David [GB/GB]; Pfizer Central Research, Ramsgate Road, Sandwich, Kent CT13 9NJ (GB). MAGEE, Thomas, Victor [US/US]; Pfizer Central Research, Eastern Point Road, Groton, CT 06340 (US). MAW, Graham, Nigel [GB/GB]; Pfizer Central Research, Ramsgate Road, Sandwich, Kent CT13 9NJ (GB). MIDDLETON, Donald, Stuart [GB/GB]; Pfizer Central Research, Ramsgate Road, Sandwich, Kent CT13 9NJ (GB). (74) Agents: RUDDOCK, Keith, Stephen et al.; Pfizer Limited, Ramsgate Road, Sandwich, Kent CT13 9NJ (GB).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: AZETIDINYLPROPYLPIPERIDINE DERIVATIVES, INTERMEDIATES AND USE AS TACHYKININ ANTAGONISTS		
(57) Abstract Compounds of formula (I) and pharmaceutically acceptable salts thereof are useful as tachykinin inhibitors which act at the NK ₁ , NK ₂ and NK ₃ receptors or a combination of two or more thereof. In said formula (I) A is CO, SO ₂ or CH ₂ ; Ar ¹ is phenyl optionally substituted by one or more substituents independently selected from halogen, C ₁₋₆ alkoxy optionally substituted by one or more halogen, and C ₁₋₆ alkyl optionally substituted by one or more halogen; X ¹ is C ₃₋₇ cycloalkyl, aryl or C ₁₋₆ alkyl, said C ₁₋₆ alkyl being optionally substituted by fluoro, CO ₂ H, CO ₂ (C ₁₋₄ alkyl), C ₃₋₇ cycloalkyl, adamantyl, aryl or het, and said C ₃₋₇ cycloalkyl being optionally substituted by 1 or 2 substituents each independently selected from C ₁₋₄ alkyl, C ₃₋₇ cycloalkyl, C ₁₋₄ alkoxy, OH, F, fluoro(C ₁₋₄ alkyl) and fluoro(C ₁₋₄ alkoxy); X is a direct link or NR ¹ ; R is SO ₂ aryl, SO ₂ (C ₁₋₆ alkyl optionally substituted by one or more halogen), (a) or (b). <div style="text-align: center; margin-top: 20px;"> <div style="display: flex; justify-content: space-around; margin-top: 10px;"> <div style="text-align: center;"> <p>(a)</p> </div> <div style="text-align: center;"> <p>(b)</p> </div> </div> </div>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

AZETIDINYLPROPYLPYPERIDINE DERIVATIVES, INTERMEDIATES AND USE AS TACHYKININ ANTAGONISTS

This invention relates to therapeutic agents of the azetidine family. More particularly, this invention relates to azetidinypropylpiperidine derivatives and to processes for the preparation of, intermediates used in the preparation of, compositions containing and uses of, such derivatives.

International Patent Application Publication Number WO 96/05193 discloses various (azetidin-1-ylalkyl)lactams as tachykinin antagonists.


International Patent Application Publication Number WO97/25322 discloses various azetidinyalkyl derivatives of N-substituted nitrogen heterocycles as tachykinin antagonists. Certain of the compounds and salts of this invention are generically but not specifically described therein.

The present azetidinypropylpiperidine derivatives are antagonists of tachykinins, including neurokinin A (NKA), neurokinin B (NKB) and Substance P, acting at the human neurokinin-1 (NK₁), neurokinin-2 (NK₂) or neurokinin-3 (NK₃) receptor, or a combination of two or more thereof. They are therefore useful for preventing or treating an inflammatory disease such as arthritis, psoriasis, asthma or inflammatory bowel disease, a central nervous system (CNS) disorder such as anxiety, depression, dementia or psychosis, a gastro-intestinal (GI) disorder such as functional bowel disease, dyspepsia, irritable bowel syndrome, gastro-oesophageal reflux, faecal incontinence, colitis, ulcerative colitis or Crohn's disease, a disease caused by Helicobacter pylori or other urease positive Gram negative bacteria, a urogenital tract disorder such as incontinence, hyperreflexia, impotence or cystitis, a pulmonary disorder such as chronic obstructive airways disease, an allergy such as eczema, contact dermatitis, atopic dermatitis, urticaria, eczematoid dermatitis or rhinitis, a hypersensitivity disorder such as to poison ivy, a vasospastic disease such as angina or Reynaud's disease, a proliferative disorder such as cancer or a disorder involving fibroblast proliferation, a fibrosing or collagen disease such as scleroderma or eosinophilic fasciitis, reflux sympathetic dystrophy such as shoulder/hand syndrome, an addiction disorder such as alcoholism, a stress-related somatic disorder, a peripheral neuropathy such as diabetic neuropathy, neuralgia, causalgia, painful neuropathy, a burn, herpetic neuralgia or post-herpetic neuralgia, a neuropathological disorder

such as Parkinson's disease, Alzheimer's disease or multiple sclerosis, a disorder related to immune enhancement or suppression such as systemic lupus erythematosus, a rheumatic disease such as fibrositis, emesis, cough, acute or chronic pain, migraine, an ophthalmic disease such as proliferative retinopathy, ocular inflammation, conjunctivitis, a bladder disorder, or a viral disease such as influenza or a cold.

The present derivatives are particularly potent and selective antagonists of tachykinins, including NKA, NKB and Substance P, acting at the human NK₁, NK₂ and NK₃ receptors or combinations of two or more thereof. They are particularly useful for treating or preventing an inflammatory disease such as arthritis, psoriasis, asthma or inflammatory bowel disease, a central nervous system (CNS) disorder such as anxiety, depression, dementia or psychosis, a gastro-intestinal (GI) disorder such as functional bowel disease, dyspepsia, irritable bowel syndrome, gastro-oesophageal reflux, faecal incontinence, colitis, ulcerative colitis or Crohn's disease, a urogenital tract disorder such as incontinence or cystitis, a neuropathological disorder such as Parkinson's disease, Alzheimer's disease or multiple sclerosis, a pulmonary disorder such as chronic obstructive airways disease, an allergy such as eczema, contact dermatitis or rhinitis, a hypersensitivity disorder such as to poison ivy, a peripheral neuropathy such as diabetic neuropathy, neuralgia, causalgia, painful neuropathy, ocular inflammation, conjunctivitis, a bladder disorder, a burn, herpetic neuralgia or post-herpetic neuralgia, cough or acute or chronic pain.

The present invention provides compounds of the formula (I):



and the pharmaceutically-acceptable salts thereof,

wherein

A is CO, SO₂ or CH₂;

Ar¹ is phenyl optionally substituted by one or more substituents independently selected from halogen, C₁₋₆ alkoxy optionally substituted by one or more halogen, and C₁₋₆ alkyl optionally substituted by one or more halogen;


X¹ is C₃₋₇ cycloalkyl, aryl or C₁₋₆ alkyl,

said C₁₋₆ alkyl being optionally substituted by fluoro, CO₂H, CO₂(C₁₋₄ alkyl), C₃₋₇ cycloalkyl, adamantyl, aryl or het,


and said C₃₋₇ cycloalkyl being optionally substituted by 1 or 2 substituents each independently selected from C₁₋₄ alkyl, C₃₋₇ cycloalkyl, C₁₋₄ alkoxy, OH, F, fluoro(C₁₋₄ alkyl) and fluoro(C₁₋₄ alkoxy);

X is a direct link or NR¹;

R is SO₂aryl, SO₂(C₁₋₆ alkyl optionally substituted by one or more halogen),



or




R¹ is H or C₁₋₆ alkyl optionally substituted by one or more halogen,


or R¹ is phenyl optionally substituted by one or more substituents independently selected from halogen, C₁₋₆ alkoxy optionally substituted by one or more halogen, or C₁₋₆ alkyl optionally substituted by one or more halogen;

Y is O, NCO(C₁₋₆ alkyl optionally substituted by one or more halogen), NCO₂(C₁₋₆ alkyl optionally substituted by one or more halogen), NSO₂(C₁₋₆ alkyl optionally substituted by one or more halogen), NCOaryl, NCO₂aryl, NSO₂aryl, NSO₂(C₁₋₆ alkyl optionally substituted by one or more halogen), CH₂, CHF, CF₂, NH, NCH₂aryl, N(C₁₋₆ alkyl optionally substituted by one or more halogen), or NCH₂(C₃₋₇ cycloalkyl);

with the proviso that X is not NR^1 when R is



or



wherein "aryl" used in all the above definitions represents phenyl or naphthyl each optionally substituted with one or more substituents independently selected from halogen, C_{1-6} alkoxy optionally substituted by one or more halogen, and C_{1-6} alkyl optionally substituted by one or more halogen;

and "het" used in the definition of X^1 means thienyl or a 5- or 6-membered ring heteroaryl group containing either 1 or 2 nitrogen heteroatoms or 1 nitrogen heteroatom and one oxygen or sulphur heteroatom, each optionally substituted by 1 or 2 substituents each independently selected from C_{1-4} alkyl, C_{1-4} alkoxy, halogen, fluoro(C_{1-4} alkyl) and fluoro(C_{1-4} alkoxy),

hereinafter referred to as "substances of the invention".

"Alkyl" groups can be straight or branched chain.

"Halogen" means fluorine, chlorine, bromine or iodine.

"Ph" means phenyl.

The pharmaceutically acceptable salts of the compounds of the formula (I) include the acid addition and the base salts thereof.

Suitable acid addition salts are formed from acids which form non-toxic salts and examples are the hydrochloride, hydrobromide, hydroiodide, sulphate, hydrogen sulphate, nitrate, phosphate,

hydrogen phosphate, acetate, maleate, fumarate, lactate, tartrate, citrate, gluconate, succinate, benzoate, methanesulphonate, benzenesulphonate and p-toluenesulphonate salts.

Suitable base salts are formed from bases which form non-toxic salts and examples are the aluminium, calcium, lithium, magnesium, potassium, sodium, zinc and diethanolamine salts.

For a review on suitable salts see Berge *et al*, J. Pharm. Sci., **66**, 1-19 (1977).

Preferably, A is CO or SO₂.

More preferably, A is CO.

Preferably, Ar¹ is phenyl optionally substituted by one or more halogen atoms.

More preferably, Ar¹ is phenyl optionally substituted by up to two halogen atoms.

Yet more preferably, Ar¹ is phenyl optionally substituted by one or two chlorine atoms.

Most preferably, Ar¹ is phenyl, 4-chlorophenyl or 3,4-dichlorophenyl.

Preferably X¹ is aryl or C₁₋₆ alkyl,

said C₁₋₆ alkyl being optionally substituted by fluoro, CO₂H, CO₂(C₁₋₄)alkyl, C₃₋₇ cycloalkyl, adamantyl, aryl or het,

and said C₃₋₇ cycloalkyl being optionally substituted by 1 or 2 substituents each independently selected from C₁₋₄ alkyl, C₃₋₇ cycloalkyl, C₁₋₄ alkoxy, OH, F, fluoro(C₁₋₄ alkyl) and fluoro(C₁₋₄ alkoxy).

More preferably, X¹ is C₁₋₆ alkyl substituted by C₃₋₇ cycloalkyl,

said C₃₋₇ cycloalkyl being optionally substituted by 1 or 2 substituents each independently selected from C₁₋₄ alkyl, C₃₋₇ cycloalkyl, C₁₋₄ alkoxy, OH, F, fluoro(C₁₋₄ alkyl) and fluoro(C₁₋₄ alkoxy),

or X¹ is phenyl optionally substituted by one or more substituents independently selected from halogen and C₁₋₆ alkyl optionally substituted by one or more halogen.

Yet more preferably, X¹ is C₁₋₆ alkyl substituted by C₃₋₇ cycloalkyl,

or is phenyl optionally substituted by one or more substituents independently selected from halogen and C₁₋₃ alkyl optionally substituted by one or more halogen.

Even more preferably, X¹ is (C₃₋₇ cycloalkyl)methyl, or

phenyl optionally substituted by one or more substituents independently selected from halogen and methyl optionally substituted by one or more halogen.

Even yet more preferably, X^1 is cyclopropylmethyl or phenyl optionally substituted by one or more fluorine or chlorine atoms.


Most preferably, X^1 is cyclopropylmethyl, phenyl, 3-chlorophenyl or 3,4-difluorophenyl.

Preferably, X is a direct link or NR^1 , where R^1 is C_{1-6} alkyl optionally substituted by one or more halogen, or R^1 is phenyl optionally substituted by one or more substituents independently selected from halogen, C_{1-6} alkoxy optionally substituted by one or more halogen, or C_{1-6} alkyl optionally substituted by one or more halogen.


More preferably, X is a direct link or NR^1 , where R^1 is C_{1-6} alkyl optionally substituted by one or more halogen, or R^1 is phenyl optionally substituted by one or more halogen substituents.

Most preferably, X is a direct link, N-methyl or N-phenyl.

Preferably R is SO_2 aryl, $SO_2(C_{1-6}$ alkyl optionally substituted by one or more halogen),




or




where Y is O, $NCO(C_{1-6}$ alkyl optionally substituted by one or more halogen), $NCO_2(C_{1-6}$ alkyl optionally substituted by one or more halogen), $NSO_2(C_{1-6}$ alkyl optionally substituted by one or more halogen), NCO aryl, CH_2 , CHF , CF_2 , NH , NCH_2 aryl, $N(C_{1-6}$ alkyl optionally substituted by one or more halogen), or $NCH_2(C_{3-7}$ cycloalkyl).

More preferably R is SO_2 (phenyl optionally substituted with one or more substituents independently selected from halogen, C_{1-6} alkoxy optionally substituted by one or more halogen, or C_{1-6} alkyl optionally substituted by one or more halogen), $SO_2(C_{1-3}$ alkyl optionally substituted by one or more halogen),




or




where Y is O, NCO(C₁₋₆ alkyl), NCO₂(C₁₋₆ alkyl), NSO₂(C₁₋₆ alkyl optionally substituted by one or more halogen), NCO(phenyl optionally substituted with one or more substituents independently selected from halogen, C₁₋₆ alkoxy optionally substituted by one or more halogen, or C₁₋₆ alkyl optionally substituted by one or more halogen), CHF, CF₂, NH, NCH₂(phenyl optionally substituted with one or more substituents independently selected from halogen, C₁₋₆ alkoxy optionally substituted by one or more halogen, or C₁₋₆ alkyl optionally substituted by one or more halogen), N(C₁₋₆ alkyl), or NCH₂(C₃₋₅ cycloalkyl).

Yet more preferably R is SO₂Ph, SO₂CH₃,




or




where Y is O, NCOCH₃, NCO₂C(CH₃)₃, NSO₂CH₃, NSO₂C₂H₅, NSO₂Ph, NSO₂CH₂Ph, NSO₂CH(CH₃)₂, NSO₂CH₂CF₃, NCOPh, CHF, NH, NCH₂Ph, or N-cyclopropylmethyl.

Most preferably R is SO₂Ph, SO₂CH₃,



or



where Y is O, NCOCH₃, NCO₂C(CH₃)₃, NSO₂CH₃, NSO₂C₂H₅, NSO₂Ph, NSO₂CH₂Ph, NSO₂CH(CH₃)₂, NSO₂CH₂CF₃, NCOPh, or CHF.

The most preferred substances of the invention are the compounds mentioned in the Examples below and the salts thereof.

Certain of the compounds of the invention can exist in one or more stereoisomeric forms. The present invention includes all such individual isomers and salts thereof.

Separation of diastereomers may be achieved by conventional techniques, e.g. by fractional crystallisation, chromatography or H.P.L.C. of a stereoisomeric mixture of a compound of formula (I) or a salt thereof. An individual enantiomer of a compound of formula (I) may also be prepared from a corresponding optically pure intermediate or by resolution, such as by H.P.L.C. of the corresponding racemate using a suitable chiral support or by fractional crystallisation of the diastereomeric salts formed by reaction of the corresponding racemate with a suitably optically active base or acid.



The invention further provides synthetic methods for the production of the substances of the invention, which are described below and in the Examples. The skilled man will appreciate that the compounds of the invention could be made by methods other than those herein described, by adaptation of the methods herein described and/or adaptation of methods known in the art, for example the art described herein.

In the Methods below, unless otherwise specified, the substituents are as defined above with reference to the compounds of formula (I) above.

Synthetic Methods

Method 1

The compounds of the invention may be prepared by reaction of a compound of formula (II)




where Z is a suitable leaving group such as methanesulphonate, toluenesulphonate, Cl, Br or I, with a compound of formula (III), optionally in the presence of a base, such as potassium carbonate, preferably in an inert solvent, such as acetonitrile.

The compounds of formulae (II) and (III) may be made by conventional methods, for example by the methods described in the Examples and Preparations, and by adaptation thereof.


Method 2

The compounds of the invention may be prepared by reaction of a compound of formula (IV) with a compound of the formula (III) as defined in Method 1 above, or a salt thereof. The reductive amination reaction is preferably carried out in the presence of a suitable acid, such as acetic acid.



(IV)

The reaction proceeds via the initial formation of an intermediate iminium salt of the formula (V), where Y^- is the counterion of the acid HY , which salt of the formula (V) may be stable and isolatable.



(V)

The reaction is preferably carried out without isolation of the intermediate of formula (V) in which case it is reduced *in situ* to provide a compound of formula (I). In a typical procedure,


the aldehyde of formula (IV) is first reacted with the azetidine of formula (III) or salt thereof in a suitable solvent, such as tetrahydrofuran (THF), and the mixture is then treated with a suitable reducing agent, e.g. sodium triacetoxyborohydride or sodium cyanoborohydride, in the presence of a suitable acid, e.g. acetic acid, to give the required product. If an acid addition salt of the azetidine of formula (III) is used as starting material, a suitable acid acceptor, e.g. triethylamine can be added prior to the addition of the reducing agent.

The reaction is typically carried out at room temperature.

The compounds of formula (IV) may be made by oxidation of compounds of formula (II) as described in Method 1 above, where Z is OH. Suitable oxidising systems include the "Swern" system, using oxalyl chloride/methyl sulphoxide in the presence of a base such as diisopropylethylamine.

Method 3

Compounds of formula (I) can be made by reaction of compounds of formula (VI) with compounds of formula (VII), where Z^1 is a suitable leaving group such as is defined above for Z in Method 1.



Where A is CO, Z^1 is conveniently Cl, or $OCOX^1$ (i.e. the reagent is the acid anhydride of X^1CO_2H). The acylation can be carried out using conventional methods, such as in the presence of a suitable base in a suitable solvent.


Alternatively, compounds where A is CO can be conveniently prepared by reaction of a compound of formula (VI) with an acid of formula (VII) where Z^1 is OH, in a "peptide coupling" reaction, such as is described by RC Larock in "Comprehensive Organic Transformations - A Guide to Functional Group Preparations", VCH (1989), and references therein.

Where A is CH₂, the alkylation can be carried out by conventional methods, such as by reaction of the compound of formula (VI) with an species of formula (VII) where Z¹ is a suitable leaving group such as Cl, Br, I, methanesulphonate or toluenesulphonate, optionally in the presence of a suitable base such as triethylamine.

Where A is SO₂, the reaction can suitably be carried out with a species of formula (VII) where Z¹ is a suitable leaving group such as Cl, Br, I, methanesulphonate or toluenesulphonate, optionally in the presence of a suitable base such as triethylamine.

Method 4

Piperidine compounds of formula (I) can be prepared by reduction of the corresponding lactams of formula (VIII), for example by reaction with a suitable hydride reducing agent such as are described in RC Larock in "Comprehensive Organic Transformations - A Guide to Functional Group Preparations", VCH (1989), pp.432-434, and references therein.




(VIII)

Compounds of formula (VIII) can be made by methods described in WO 96/05193 and adaptation thereof by methods known in the art.

Method 5

Compounds of formula (I) can be prepared by reaction of compounds of formula (IX), where Z² is a leaving group and corresponds to Z as defined in Method 1 above, with a nucleophilic reagent acting as a RX⁻ "synthon".




(IX)

Suitably, Z^2 is methanesulphonate, and the reaction is carried out by generation of the RX^- moiety by e.g. reaction of the corresponding RXH compound with a base such as an alkyllithium base, optionally in the presence of an inert solvent. Alternatively the RXH compound may itself be nucleophilic enough to displace the Z^2 group without first converting it to the anion RX^- .

Compounds of formula (IX) can be prepared by adaptation of methods described herein and in the art described herein, by methods known to the skilled chemist.

Method 6

Compounds of formula (I) may be prepared from compounds of formula (X) where Z^3 and Z^4 are leaving groups independently defined as Z in Method 1, with a reagent of formula X^1-A-NH_2 .




(X)


Compounds of formulae (X) and X^1-A-NH_2 may be prepared by conventional methods, such as by adaptation of methods described herein, and in the art described herein.

Method 7


Compounds of the formula (I) where R is




or



where Y is NCO(C₁₋₆ alkyl optionally substituted by one or more halogen), NCO₂(C₁₋₆ alkyl optionally substituted by one or more halogen), NSO₂(C₁₋₆ alkyl optionally substituted by one or more halogen), NCOaryl, NCO₂aryl, NSO₂aryl, NSO₂(C₁₋₆ alkyl optionally substituted by one or more halogen), NCH₂aryl, N(C₁₋₆ alkyl optionally substituted by one or more halogen), or NCH₂(C₃₋₇ cycloalkyl), can be made by reaction of a compound of formula (I) where R is




or




as appropriate, and Y is NH, with a reagent of formula [(C₁₋₆ alkyl optionally substituted by one or more halogen)CO-, (C₁₋₆ alkyl optionally substituted by one or more halogen)CO₂-, (C₁₋₆ alkyl optionally substituted by one or more halogen)SO₂-, arylCO-, arylCO₂-, arylSO₂-, (C₁₋₆ alkyl optionally substituted by one or more halogen)SO₂-, arylCH₂-, (C₁₋₆ alkyl optionally substituted by one or more halogen)-, or (C₃₋₇ cycloalkyl)CH₂-]-Z⁵, where Z⁵ is a suitable leaving group such as chloride, bromide, iodide, methanesulphonate, toluenesulphonate, etc., optionally in the presence of a base such as triethylamine, preferably in the presence of an inert solvent.

Method 8

Compounds of the formula (I) where R is



or



where Y is NH can be conveniently prepared by reaction of a corresponding compound where Y is $\text{NCO}_2\text{C}(\text{CH}_3)_3$, with an acid, such as trifluoroacetic acid, preferably in the presence of an inert solvent.

Where desired or necessary, the compound of formula (I) can be converted into a pharmaceutically acceptable salt thereof, conveniently by mixing together solutions of a compound of formula (I) and the desired acid or base, as appropriate. the salt may be precipitated from solution and collected by filtration, or may be collected by other means such as by evaporation of the solvent.

It is to be understood that the synthetic transformation methods mentioned herein may be carried out in various different sequences in order that the desired compounds can be efficiently assembled. The skilled chemist will exercise his judgement and skill as to the most efficient sequence of reactions for synthesis of a given target compound.

It will be apparent to those skilled in the art that sensitive functional groups may need to be protected and deprotected during synthesis of a compound of the invention. This may be achieved by conventional methods, for example as described in "Protective Groups in Organic Synthesis" by TW Greene and PGM Wuts, John Wiley & Sons Inc (1991).

The compounds of the invention may be purified by conventional methods.

The affinity of the compounds of formula (I) and their salts for the human NK_1 receptor can be tested in vitro by testing their ability to inhibit [^3H]-Substance P binding to membranes prepared from the human IM9 cell line expressing the human NK_1 receptor using a modification of the method described in McLean, S. et al, J. Pharm. Exp. Ther., 267, 472-9 (1993) in which whole cells were used.

The affinity of the compounds of formula (I) and their salts for the human NK_2 receptor can be tested in vitro by testing their ability to compete with [^3H] or [^{125}I]NKA (neurokinin A) for binding to membranes prepared from Chinese hamster ovary cells expressing the cloned

human NK₂ receptor. In this method, washed Chinese hamster ovary cell membranes are prepared as described for the previous method where IM9 cells are used instead. The membranes are incubated (90 min, 25°C) with [³H] or [¹²⁵I] NKA and with a range of concentrations of the test compound. Non-specific binding was determined in the presence of 10μM NKA.

The NK₂ receptor antagonist activity of the compounds of the formula (I) can be tested, in vitro, by testing their ability to antagonise the contractile effects of the selective NK₂ receptor agonist [βAla⁸]NKA₍₄₋₁₀₎ in the rabbit pulmonary artery, using the method of Patacchini and Maggi, Eur. J. Pharmacol., 236, 31-37 (1993).

The compounds of the formula (I) and their salts can be tested for NK₂ receptor antagonist activity, in vivo, by testing their ability to inhibit bronchoconstriction induced by [βAla⁸]NKA₍₄₋₁₀₎ in the anaesthetised guinea pig, using the method described by Murai et al, J. Pharm. Exp. Ther., 262, 403-408 (1992) or Metcalfe et al, Br. J. Pharmacol., 112, 563P (1994).

The compounds of the formula (I) and their salts can be tested for NK₃ receptor antagonist activity, in vitro, by testing their ability to antagonise the contractile effects of the selective NK₃ receptor agonist senktide in the guinea-pig ileum using the method of Maggi et al, Br. J. Pharmacol., 101, 996-1000 (1990).

The affinity of the compounds of formula (I) and their salts for the NK₃ receptor can be tested in vitro by testing their ability to displace [³H] senktide at the guinea pig cortex NK₃ receptor. The receptors are prepared by the method described by S.Guard, et al, in Br.J.Pharmacol. (1990), 99, 767.

For human use, the compounds of the formula (I) and their salts can be administered alone, but will generally be administered in admixture with a pharmaceutically acceptable diluent or carrier selected with regard to the intended route of administration and standard pharmaceutical practice. For example, they can be administered orally, including sublingually, in the form of tablets containing such excipients as starch or lactose, or in capsules or ovules either alone or in admixture with excipients, or in the form of elixirs, solutions or suspensions containing flavouring or colouring agents. They can be injected parenterally, for example,

intravenously, intramuscularly or subcutaneously. For parenteral administration, they are best used in the form of a sterile aqueous solution which may contain other substances, for example, enough salts or glucose to make the solution isotonic with blood.

For oral and parenteral administration to human patients, the daily dosage level of the compounds of the formula (I) and their salts will be from 0.001 to 20, preferably from 0.01 to 20, more preferably from 0.1 to 10, and most preferably from 0.5 to 5, mg/kg (in single or divided doses). Thus tablets or capsules of the compounds will contain from 0.1 to 500, preferably from 50 to 200, mg of active compound for administration singly or two or more at a time, as appropriate. The physician in any event will determine the actual dosage which will be most suitable for an individual patient and it will vary with the age, weight and response of the particular patient. The above dosages are exemplary of the average case; there can, of course, be individual instances where higher or lower dosage ranges are merited, and such are within the scope of this invention.

Alternatively, the compounds of the formula (I) can be administered by inhalation or in the form of a suppository or pessary, or they may be applied topically in the form of a lotion, solution, cream, ointment or dusting powder. An alternative means of transdermal administration is by use of a skin patch. For example, they can be incorporated into a cream consisting of an aqueous emulsion of polyethylene glycols or liquid paraffin; or they can be incorporated, at a concentration between 1 and 10%, into an ointment consisting of a white wax or white soft paraffin base together with such stabilisers and preservatives as may be required.

It is to be appreciated that reference to treatment includes prophylaxis as well as the alleviation of established symptoms of the disease.

The invention further provides:-

- i) a pharmaceutical composition comprising a substance of the invention together with a pharmaceutically acceptable diluent or carrier;
- ii) a substance of the invention or composition thereof, for use as a medicament;

- iii) the use of a substance of the invention or composition thereof, for the manufacture of a medicament for the treatment of a disease by producing an antagonist effect on a tachykinin acting at the human NK₁, NK₂ or NK₃ receptor, or a combination of two or more thereof;
- iv) use as in (iii) where the disease is an inflammatory disease such as arthritis, psoriasis, asthma or inflammatory bowel disease, a central nervous system (CNS) disorder such as anxiety, depression, dementia or psychosis, a gastro-intestinal (GI) disorder such as functional bowel disease, dyspepsia, irritable bowel syndrome, gastro-oesophageal reflux, faecal incontinence, colitis, ulcerative colitis or Crohn's disease, a disease caused by Helicobacter pylori or other urease positive Gram negative bacteria, a urogenital tract disorder such as incontinence, hyperreflexia, impotence or cystitis, a pulmonary disorder such as chronic obstructive airways disease, an allergy such as eczema, contact dermatitis, atopic dermatitis, urticaria, eczematoid dermatitis or rhinitis, a hypersensitivity disorder such as to poison ivy, a vasospastic disease such as angina or Reynaud's disease, a proliferative disorder such as cancer or a disorder involving fibroblast proliferation, a fibrosing or collagen disease such as scleroderma or eosinophilic fasciitis, reflux sympathetic dystrophy such as shoulder/hand syndrome, an addiction disorder such as alcoholism, a stress-related somatic disorder, a peripheral neuropathy such as diabetic neuropathy, neuralgia, causalgia, painful neuropathy, a burn, herpetic neuralgia or post-herpetic neuralgia, a neuropathological disorder such as Parkinson's disease, Alzheimer's disease or multiple sclerosis, a disorder related to immune enhancement or suppression such as systemic lupus erythematosus, a rheumatic disease such as fibrositis, emesis, cough, acute or chronic pain, migraine, an opthalmic disease such as proliferative retinopathy, ocular inflammation, conjunctivitis, a bladder disorder, or a viral disease such as influenza or a cold;
- v) a method of treatment of a human to treat a disease by producing an antagonist effect on a tachykinin acting at the human NK₁, NK₂ or NK₃ receptor, or a combination of two or more thereof, which comprises treating said human with an effective amount of a substance of the invention or composition thereof;
- vi) a method as in (v) where the disease is an inflammatory disease such as arthritis, psoriasis, asthma or inflammatory bowel disease, a central nervous system (CNS) disorder such as anxiety, depression, dementia or psychosis, a gastro-intestinal (GI)

disorder such as functional bowel disease, dyspepsia, irritable bowel syndrome, gastro-oesophageal reflux, faecal incontinence, colitis, ulcerative colitis or Crohn's disease, a disease caused by Helicobacter pylori or other urease positive Gram negative bacteria, a urogenital tract disorder such as incontinence, hyperreflexia, impotence or cystitis, a pulmonary disorder such as chronic obstructive airways disease, an allergy such as eczema, contact dermatitis, atopic dermatitis, urticaria, eczematoid dermatitis or rhinitis, a hypersensitivity disorder such as to poison ivy, a vasospastic disease such as angina or Reynaud's disease, a proliferative disorder such as cancer or a disorder involving fibroblast proliferation, a fibrosing or collagen disease such as scleroderma or eosinophilic fasciitis, reflux sympathetic dystrophy such as shoulder/hand syndrome, an addiction disorder such as alcoholism, a stress-related somatic disorder, a peripheral neuropathy such as diabetic neuropathy, neuralgia, causalgia, painful neuropathy, a burn, herpetic neuralgia or post-herpetic neuralgia, a neuropathological disorder such as Parkinson's disease, Alzheimer's disease or multiple sclerosis, a disorder related to immune enhancement or suppression such as systemic lupus erythematosus, a rheumatic disease such as fibrositis, emesis, cough, acute or chronic pain, migraine, an ophthalmic disease such as proliferative retinopathy, ocular inflammation, conjunctivitis, a bladder disorder, or a viral disease such as influenza or a cold;

- vii) the synthetic methods herein described,
- and
- viii) certain novel intermediates herein described.


The following Examples illustrate the preparation of the substances of the invention.

In the Examples and Preparations which follow, ^1H NMR data are presented in parts per million shift from tetramethylsilane (δ), and are measured in solution in CDCl_3 unless specified otherwise. Thin layer chromatography (TLC) was carried out using Merck kieselgelTM silica plates, unless otherwise specified.

In the structures which follow "Me" and "Ph" represent a methyl and phenyl group respectively.

EXAMPLE 1

(+)-3(R)-1-Benzoyl-3-(3-[3-(4-*tert*-butoxycarbonyl)piperazin-1-yl]azetidin-1-yl]propyl)-3-(3,4-dichlorophenyl)piperidine.



A mixture of the compounds of PREPARATION 8 (11.62g) and PREPARATION 10 (14g) and potassium carbonate (10.23g) in dry acetonitrile (300mL) was heated, with vigorous stirring, under reflux for 18 hours. The cooled mixture was diluted with ethyl acetate (700mL) and washed with 50% aqueous sodium chloride solution (70mL). The organic solution was dried over sodium sulphate and evaporated to dryness *in vacuo*. The residue was purified by column chromatography over silica gel using gradient elution (dichloromethane/methanol/ammonium hydroxide 93/7/1 to 80/20/3) to afford the title compound (15.36g) and recovered compound from PREPARATION 10 (5.85g).

TLC R_f : 0.15 (dichloromethane/methanol 95/5 by volume).

LRMS m/z : 616 (MH^+).

1H NMR : 7.38 (m, 4H), 7.21 (m, 4H), 4.52 (br s, 1H), 3.38 (m, 9H), 2.90 (m, 1H), 2.72 (m, 2H), 2.37-2.03 (m, 8H), 1.80 (m, 1H), 1.60 (m, 2H), 1.42 (s, 9H), 1.20 (m, 1H), 0.93 (m, 2H).

Found: C, 63.32; H, 7.21; N, 9.05. $C_{33}H_{44}Cl_2N_4O_3 \cdot 3/20CH_2Cl_2$ requires C, 63.36; H, 7.11; N, 8.92%.

$[a]_D +16.2^\circ$ ($c=0.13$, methanol).

EXAMPLE 2

3(R)-1-Benzoyl 3-(3-[3-(4-*tert*-butoxycarbonyl)homopiperazin-1-yl]azetidin-1-yl]propyl)-3-(3,4-dichlorophenyl)piperidine


The compounds from PREPARATION 8 (470 mg) and PREPARATION 12 (260 mg) were mixed with potassium carbonate (414 mg) in acetonitrile (15 mL) and heated to reflux overnight. The reaction mixture was allowed to cool and was partitioned between ethyl acetate and water. The organic phase was washed with brine, dried over magnesium sulphate, and filtered to give a crude material which was subsequently purified by silica gel chromatography with dichloromethane:methanol (98:2) eluant to give pure product (334 mg).

^1H NMR : 7.35-7.2 (m, 8H), 3.45-3.37 (m, 13H), 2.59-2.06 (m, 6H), 1.74-1.40 (m, 10H).

LRMS m/z 630 ($m+H$)⁺

EXAMPLE 3

(+)-3(R)-1-Benzoyl-3-(3,4-dichlorophenyl)-3-(3-[3-(4-methanesulphonylpiperazin-1-yl)azetidin-1-yl]propyl)piperidine



A solution of methanesulphonyl chloride (122 μ l) in dichloromethane (15mL) was added over a period of 30 minutes to an ice-cold solution of the compound of PREPARATION 13 (773mg) and triethylamine (420 μ l) in dichloromethane (30mL) under nitrogen and the mixture stirred for 1 hour. The reaction mixture was evaporated to dryness *in vacuo* and the residue was purified by column chromatography over silica gel (dichloromethane/methanol/ammonium hydroxide 95/5/0.5) to afford the title compound as a foam (740mg).

TLC R_f : 0.41 (dichloromethane/methanol/ammonium hydroxide 93/7/1 by volume).

m/z : 593 (MH^+).


^1H NMR : 7.38 (m, 5H), 7.20 (m, 3H), 4.48 (m, 1H), 3.60-3.08 (m, 8H), 2.92 (m, 1H), 2.70 (m, 5H), 2.30 (m, 6H), 2.10 (br s, 1H), 1.78 (m, 1H), 1.68-1.05 (m, 6H), 0.90 (m, 1H).

Found: C, 56.77; H, 6.19; N, 9.03. $C_{29}H_{38}Cl_2N_4O_3S \cdot 3/10CH_2Cl_2$ requires C, 56.84; H, 6.28; N, 9.05%.

$[\alpha]_D +20.3^\circ$ ($c=0.12$, methanol).

EXAMPLE 4

(+)-3(R)-3-(3-[3-(4-Acetyl)piperazin-1-yl]azetidin-1-yl]propyl)-1-benzoyl-3-(3,4-dichlorophenyl)piperidine.



A solution of acetic anhydride (104 μ l) in dichloromethane (20mL) was added to an ice-cold solution of the compound of PREPARATION 13 (515mg) and triethylamine (280 μ l) in dichloromethane (25mL). The reaction mixture was allowed to warm to room temperature and stirring was continued for 2 hours. The reaction mixture was evaporated to dryness *in vacuo* and the residue purified by column chromatography over silica gel using gradient elution (dichloromethane/methanol/ammonium hydroxide 95/5/0.5 to 93/7/1 by volume) to afford the title compound as a foam (458mg).

TLC R_f: 0.33 (dichloromethane/methanol/ammonium hydroxide 93/7/1 by volume).

m/z: 556 (MH⁺).


¹H NMR: 7.50-7.00 (m, 8H), 4.40 (br s, 1H), 3.55 (m, 3H), 3.43-3.00 (m, 8H), 2.85 (m, 1H), 2.70 (br s, 2H), 2.37-1.82 (m, 6H), 1.82-1.01 (m, 8H), 0.88 (m, 1H).

Found: C, 62.75; H, 6.93; N, 9.61. C₃₀H₃₈Cl₂N₄O₂.1/4CH₂Cl₂ requires C, 62.75; H, 6.93; N, 9.68%.

[α]_D +20.4° (c=0.102, methanol).

EXAMPLE 5

(+)-3(R)-1-Benzoyl-3-(3-[3-(4-Benzoyl)piperazin-1-yl]azetidin-1-yl]propyl)-3-(3,4-dichlorophenyl)piperidine.



A solution of benzoyl chloride (128 μ l) in dichloromethane (15mL) was added to a solution of the compound of PREPARATION 13 (516mg) and triethylamine (280 μ l) in dichloromethane (25mL). The reaction mixture was stirred for 2 hours at room temperature. The reaction mixture was evaporated to dryness *in vacuo* and the residue was purified by column chromatography over silica gel (dichloromethane/methanol/ammonium hydroxide 95/5/0.5 by volume) to afford the title compound as a foam (489mg).

TLC R_f: 0.49 (dichloromethane/methanol/ammonium hydroxide 93/7/1 by volume).

m/z: 618 (MH⁺).


¹H NMR : 7.48-7.13 (m, 13H), 4.43 (m, 1H), 3.73 (m, 2H), 3.45-3.05 (m, 7H), 2.90 (m, 1H), 2.72 (m, 2H), 2.39-2.01 (m, 7H), 1.78 (t, 1H), 1.67-1.05 (m, 5H), 0.90 (m, 1H).

Found: C, 66.00; H, 6.45; N, 8.69. C₃₅H₄₀Cl₂N₄O₂.1/4CH₂Cl₂ requires C, 66.06; H, 6.37; N, 8.74%.

[α]_D +15.2° (c=0.14, methanol).

EXAMPLES 6-10

The following compounds of the general formula:



or salts thereof, were prepared from the compound of Preparation 13 and the corresponding sulphonyl chlorides by similar methods to that used in Example 3.

EXAMPLE 6

(+)-3(R)-1-Benzoyl-3-(3,4-dichlorophenyl)-3-(3-[3-(4-methanesulphonylpiperazin-1-yl)azetidin-1-yl]propyl)piperidine.

EXAMPLE 7

(+)-3(R)-3-(3-[3-(4-Benzenesulphonylpiperazin-1-yl)azetidin-1-yl]propyl)-1-benzoyl-3-(3,4-dichlorophenyl)piperidine.

EXAMPLE 8

(+)-3(R)-1-Benzoyl-3-(3-[3-(4-benzylsulphonylpiperazin-1-yl)azetidin-1-yl]propyl)-3-(3,4-dichlorophenyl)piperidine.

EXAMPLE 9

(+)-3(R)-1-Benzoyl-3-(3,4-dichlorophenyl)-3-(3-[3-(4-*iso*-propylsulphonylpiperazin-1-yl)azetidin-1-yl]propyl)piperidine.

EXAMPLE 10


(+)-3(R)-1-Benzoyl-3-(3,4-dichlorophenyl)-3-(3-[3-(4-(2,2,2-trifluoroethane)sulphonylpiperazin-1-yl)azetidin-1-yl]propyl)piperidine

Example No.	R	m/z	Rf *	$[\alpha]_D$ methanol	Analysis/ ¹ H NMR
6	C ₂ H ₅	607	0.38	+17.3° c=0.102	δ_{H} : 7.35 (m, 4H), 7.20 (m, 4H), 4.49 (br s, 1H), 3.28 (m, 8H), 2.90 (m, 3H), 2.68 (m, 2H), 2.30 (m, 6H), 2.10 (m, 1H), 1.88 (t, 1H), 1.56 (m, 4H), 1.38 (m, 1H), 1.32 (t, 3H), 1.18 (m, 1H), 0.90 (m, 1H). Found: C, 58.33; H, 6.62; N, 9.05. C ₂₈ H ₄₀ Cl ₂ N ₄ O ₃ .3/20CH ₂ Cl ₂ requires C, 58.37; H, 6.55; N, 9.03%
7	Ph	654	0.47	+13.8° c=0.12	δ_{H} : 7.71 (m, 2H), 7.52 (m, 3H), 7.37 (m, 4H), 7.18 (m, 4H), 4.48 (m, 1H), 3.50-3.08 (m, 5H), 3.30 (m, 5H), 2.85 (t, 1H), 2.59 (m, 2H), 2.32 (m, 3H), 2.20 (m, 2H), 2.08 (m, 1H), 1.75 (t, 1H), 1.65-1.37 (m, 4H), 1.14 (t, 1H), 0.85 (m, 1H). Found: C, 61.45; H, 6.15; N, 8.33. C ₃₄ H ₄₀ Cl ₂ N ₄ O ₃ .1/10CH ₂ Cl ₂ requires C, 61.67; H, 6.10; N, 8.44%
8	PhCH ₂	669	0.48	+15.2° c=0.104	δ_{H} : 7.40-7.12 (m, 13H), 4.48 (m, 1H), 4.15 (s, 2H), 3.48-3.20 (m, 4H), 3.09 (m, 4H), 2.84 (m, 1H), 2.63 (m, 2H), 2.30-2.02 (m, 7H), 1.78 (t, 1H) < 1.63-1.28 (m, 5H), 1.15 (m, 1H), 0.88 (m, 1H). Found: C, 60.65; H, 6.22; N, 8.00. C ₃₃ H ₄₂ Cl ₂ N ₄ O ₃ .2/5CH ₂ Cl ₂ requires C, 60.42; H, 6.13; N, 7.06%
9	CH(CH ₃) ₂	621	0.42	+21.5° c=0.104	δ_{H} : 7.40 (m, 4H), 7.27 (m, 4H), 4.50 (m, 1H), 3.37 (m, 8H), 3.18 (m, 1H), 2.95 (m, 1H), 2.75 (m, 2H), 2.36 (m, 6H), 2.14 (m, 1H), 1.90-0.90 (m, 8H), 1.35 (d, 6H). Found: C, 58.84; H, 6.79; N, 8.74. C ₃₁ H ₄₂ Cl ₂ N ₄ O ₃ .1/5CH ₂ Cl ₂ requires C, 58.68; H, 6.69; N, 8.77%
10	CF ₃ CH ₂	661	0.44	+21.0° c=0.12,	δ_{H} : 7.48 (m, 4H), 7.17 (m, 4H), 4.48 (m, 1H), 3.67 (m, 3H), 3.30 (m, 8H), 2.91 (t, 1H), 2.68 (m, 2H), 2.30 (m, 6H), 2.10 (m, 1H), 1.79 (m, 1H), 1.58 (m, 3H), 1.38 (m, 1H), 1.18 (m, 1H), 0.90 (m, 1H). Found: C, 53.66; H, 5.57; N, 8.23. C ₃₀ H ₃₇ Cl ₂ F ₃ N ₄ O ₃ .3/10CH ₂ Cl ₂ requires C, 53.70; H, 5.58; N, 8.31%

* TLC: dichloromethane/methanol/ammonium hydroxide 93/7/1 by volume

EXAMPLE 11

(+)-3(R)-1-Benzoyl-3-(3,4-dichlorophenyl)-3-(3-[3-morpholinoazetidin-1-yl]propyl)piperidine



Part A. Oxalyl chloride (0.17 mL) was added to dichloromethane (10 mL) under nitrogen and the solution cooled in a dry ice/acetone bath before adding anhydrous methyl sulphoxide (0.18 mL) dropwise and allowing to stir for 0.75 h; to this was added the compound from PREPARATION 7 (500 mg) dissolved in dichloromethane (10 mL) dropwise. After an additional 0.75 h, diisopropylethylamine (0.67 mL) was added and the mixture allowed to warm to ambient temperature before pouring into 2 N aqueous hydrochloric acid (75 mL). The resulting solution was extracted with dichloromethane three times. The combined organic phases were washed with brine and dried over sodium sulphate before concentrating to yield the crude aldehyde intermediate as a yellow oil (LRMS m/z 390 $[m]^+$) which was taken on to the next stage without purification.

Part B. The crude aldehyde from part A was dissolved in tetrahydrofuran (25 mL) along with glacial acetic acid (0.87 mL), sodium triacetoxyborohydride (432 mg), and 3-morpholinoazetidine dihydrochloride (Preparation 25, 411 mg). After stirring the mixture at ambient temperature for 12 h, water (2 mL) was added and the solvent evaporated *in vacuo*. To the resulting crude product was added 1 N aqueous sodium hydroxide (ca. 10 mL) and this mixture was extracted with dichloromethane three times. The combined organic layers were washed with brine, dried over sodium sulphate, and concentrated *in vacuo* before purification on silica gel with dichloromethane:methanol:ammonium hydroxide (95:5:0.5) as eluant to yield 553 mg of the desired compound.

^1H NMR : 7.5-7.2 (m, 8H), 4.43 (m, 1H), 3.62 (m, 4H), 3.5-3.1 (m, 5H), 2.89 (m, 1H), 2.71 (m, 2H), 2.38-2.2 (m, 6H), 2.07 (m, 1H), 1.79 (m, 1H), 1.1-1.72 (m, 5H), 0.96 (m, 1H).


$[\alpha]_D^{25} = +22.6^\circ$ ($c = 0.1$ mg/mL in MeOH).

LRMS m/z 517 ($m+H$) $^+$.

Found: C, 63.50; H, 6.84; N, 7.81. $\text{C}_{28}\text{H}_{35}\text{N}_3\text{O}_2\text{Cl}_2 \cdot 0.18\text{CH}_2\text{Cl}_2$ requires: C, 63.65, H, 6.70, N, 7.90.

EXAMPLE 12

(+)-3(R)-1-Benzoyl-3-(3,4-dichlorophenyl)-3-[3-(3-(N-Methyl benzenesulphonamido)azetidin-1-yl)propyl]piperidine



The compound from PREPARATION 8 (225 mg) was mixed with the compound from PREPARATION 20 (250 mg) and potassium hydrogen carbonate (200 mg) in acetonitrile (15 mL) and the resulting mixture was heated to reflux overnight. After cooling the reaction to ambient temperature, water (15 mL) was added and the mixture was extracted three times with ethyl acetate. The combined organic layers were washed with brine, dried over magnesium sulphate, and concentrated *in vacuo*. Purification of the crude product on silica gel with ethyl acetate:2-propanol (4:1) as eluant yielded the title compound (165 mg).

$^1\text{H NMR}$: 7.70 (m, 1H), 7.49 (m, 2H), 7.31 (m, 2H), 7.18 (m, 3H), 4.42 (m, 1H), 3.82 (m, 1H), 3.50-3.09 (m, 5H), 2.80 (m, 2H), 2.60 (br s, 3H), 2.20 (m, 2H), 2.08 (m, 1H), 1.78 (br t, 1H), 1.70-1.08 (m, 5H), 0.85 (m, 1H).


$[\alpha]_{\text{D}}^{25} = +20.4^\circ$ ($c = 1.0$ mg/mL in MeOH).

TLC (ethyl acetate:2-propanol 4:1) $R_f = 0.25$.

LRMS m/z 601 ($m+H$) $^+$

EXAMPLE 13

(+)-3(R)-1-Benzoyl-3-(3,4-dichlorophenyl)-3-[3-(3-(N-phenyl methanesulphonamido)azetidin-1-yl)propyl]piperidine



The compound from PREPARATION 8 (300 mg) was mixed with the compound from PREPARATION 22 (400 mg) and potassium hydrogen carbonate (300 mg) in acetonitrile (15 mL) and the resulting mixture was heated to reflux overnight. After cooling the reaction to ambient temperature, water (25 mL) was added and the mixture was extracted three times with ethyl acetate. The combined organic layers were washed with brine, dried over magnesium sulphate, and concentrated *in vacuo*. Chromatography of the crude product on silica gel with ethyl acetate:2-propanol (4:1) as eluant yielded the title compound (185 mg).

^1H NMR: 7.39-7.28 (m, 8H), 7.22-7.17 (m, 5H), 4.46 (m, 2H), 3.55-3.08 (m, 5H), 2.76 (s, 3H), 2.68-2.58 (m, 2H), 2.20-2.02 (m, 3H), 1.72 (m, 1H), 1.60-1.28 (m, 4H), 1.19-1.00 (m, 1H), 0.83 (m, 1H).

$[\alpha]_{\text{D}}^{25} = +13.8^\circ$ (c = 1.0 mg/mL in MeOH).

TLC (ethyl acetate:2-propanol 4:1) $R_f = 0.20$


LRMS m/z 601 (m+H) $^+$

Found: C, 61.05; H, 5.92; N, 6.87.

$\text{C}_{31}\text{H}_{35}\text{N}_3\text{O}_3\text{Cl}_2\text{S} \cdot 0.5\text{H}_2\text{O}$ requires: C, 61.08; H, 5.95; N, 6.89

EXAMPLE 14

3(R)-1-Benzoyl-3-(3,4-dichlorophenyl)-3-(3-[3-(4-methanesulphonylhomopiperazin-1-yl)azetidin-1-yl]propyl)-piperidine



The compound from PREPARATION 14 (100 mg) was dissolved in dichloromethane (2 mL) and cooled in an ice/water bath. To this were added methanesulphonyl chloride (0.02 mL) and triethylamine (0.032 mL). The reaction mixture was allowed to warm to ambient temperature

and was stirred overnight before quenching with water. The organic phase was separated, washed with brine, dried with magnesium sulphate, filtered, and concentrated. Silica gel chromatography using dichloromethane:methanol (9:1) eluant yielded the title compound (28 mg).

^1H NMR : 7.48-7.22 (m, 8H), 3.4-1.2 (m, 32H).


TLC (dichloromethane:methanol = 9:1) R_f = 0.46.

LRMS m/z 608 ($m+H$) $^+$.

Found: C, 57.57; H, 6.56; N, 8.42. $\text{C}_{30}\text{H}_{40}\text{N}_4\text{O}_3\text{Cl}_2\text{S} \cdot 0.5\text{H}_2\text{O} \cdot 0.33\text{CH}_2\text{Cl}_2$ requires: C, 57.31; H, 6.45; N, 8.81.

EXAMPLE 15

3(R)-1-Benzenesulphonyl-3-(3,4-dichlorophenyl)-3-(3-[3-(4-methanesulphonylpiperazin-1-yl)azetidin-1-yl]propyl)-piperidine



The compound from PREPARATION 18 (125 mg) was dissolved in dichloromethane (2.5 mL) and cooled in an ice/water bath. To this were added triethylamine (0.046 mL) and benzenesulphonyl chloride (0.04 mL). The reaction mixture was allowed to warm to ambient temperature and stirred overnight before quenching with water. The organic phase was separated, washed with brine, dried with magnesium sulphate, filtered, and concentrated. Silica gel chromatography using dichloromethane:methanol (95:5) eluant yielded the title compound (52 mg).

^1H NMR : 7.73 (m, 2H), 7.55 (m, 1H), 7.50 (m, 2H), 7.37 (m, 1H), 7.20 (m, 1H), 3.46-0.84 (m, 30H)


TLC (dichloromethane:methanol = 9:1) R_f = 0.45

LRMS m/z 630 ($m+H$) $^+$

Found: C, 52.35; H, 6.11; N, 8.46. $\text{C}_{28}\text{H}_{38}\text{N}_4\text{O}_4\text{Cl}_2\text{S}_2 \cdot 0.2\text{H}_2\text{O}$ requires: C, 52.38; H, 5.99; N, 8.34.

EXAMPLE 16

(+)-3(R)-1-(Cyclopropylacetyl)-3-(3,4-dichlorophenyl)-3-[3-(4-methanesulphonylpiperazin-1-yl)azetidin-1-yl]propyl)piperidine



Cyclopropylacetic acid (0.2 mL) was dissolved in dichloromethane (5 mL) and cooled in an ice/water bath. To this were added N-methyl morpholine (0.1 mL), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (95 mg), and 1-hydroxybenzotriazole hydrate (50 mg). The compound from PREPARATION 18 (150 mg) was then added after one hour, and the mixture stirred overnight at ambient temperature. The solvent was then evaporated in vacuo, and the mixture partitioned between ethyl acetate and water. The resulting organic phase was washed with brine, dried with magnesium sulphate, filtered, and concentrated. Silica gel chromatography using dichloromethane:methanol (95:5) eluant yielded the title compound.

$^1\text{H-NMR}$: 7.36-7.13 (m, 3H), 3.50-0.79 (m, 37H)

LRMS m/z 572 ($m+H$)⁺


$[\alpha]_D^{25} = +42.21^\circ$ ($c = 1.0$ mg/mL in MeOH)

Found: C, 55.33; H, 7.05; N, 9.47. $\text{C}_{27}\text{H}_{40}\text{N}_4\text{O}_3\text{S}\cdot 0.25\text{CH}_2\text{Cl}_2$

Requires: C, 55.21, H, 6.89, N, 9.45

EXAMPLE 17

(+)-3(R)-3-(3,4-Dichlorophenyl)-1-(3,4-difluorobenzoyl)-3-[3-(4-methanesulphonylpiperazin-1-yl)azetidin-1-yl]propyl)piperidine




3,4-Difluorobenzoic acid (145 mg) was dissolved in dichloromethane (5 mL) and cooled in an ice/water bath. To this were added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide

hydrochloride (95 mg), 1-hydroxybenzotriazole hydrate (50 mg), and N-methyl morpholine (0.1 mL) and the reaction was stirred for 1 h. The compound from PREPARATION 18 (150 mg) was then added and the mixture stirred at ambient temperature overnight before removing the solvent *in vacuo*. To the resulting mixture was added 1 M sodium bicarbonate and ethyl acetate; the organic layer was separated and washed with water and brine before drying with magnesium sulphate, filtering, and concentrating to a crude mixture. Silica gel chromatography using dichloromethane:methanol (95:5) eluant yielded pure product (60 mg).
¹H NMR : 7.48-6.96 (m, 6H), 3.30-1.22 (m, 30H).
 LRMS *m/z* 630 (m+H)⁺.
 [α]_D²⁵ = +16.4 ° (c = 1.0 mg/mL in MeOH).

EXAMPLE 18


(+)-3(R)-1-(3-Chlorobenzoyl)-3-(3,4-dichlorophenyl)-3-(3-[3-(4-methanesulphonylpiperazin-1-yl)azetidin-1-yl]propyl)-piperidine



3-Chlorobenzoic acid (140 mg) was dissolved in dichloromethane (5 mL) and cooled in an ice/water bath. To this were added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (95 mg), 1-hydroxybenzotriazole hydrate (50 mg), and N-methyl morpholine (0.1 mL) and the reaction was stirred for 1 h. The compound from PREPARATION 18 (150 mg) was then added and the mixture stirred at ambient temperature overnight before removing the solvent *in vacuo*. To the resulting mixture was added 1 M sodium bicarbonate and ethyl acetate; the organic layer was separated and washed with water and brine before drying with magnesium sulphate, filtering, and concentrating to a crude mixture. Silica gel chromatography using dichloromethane:methanol (95:5) eluant yielded the title compound.
¹H-NMR : 7.42-7.06 (m, 6H), 3.38-1.37 (m, 30H)
 LRMS *m/z* 629 (m+H)⁺
 [α]_D²⁵ = +13.4 ° (c = 1.0 mg/mL in MeOH)
 Found: C, 55.17; H, 6.08; N, 8.70. C₂₉H₃₇N₄O₃SCl₃
 Requires: C, 55.46, H, 5.94, N, 8.92

EXAMPLE 19

(+)-3(R)-1-Benzoyl-3-(3,4-dichlorophenyl)-3-(3-[3-piperazinoazetidin-1-yl]propyl)piperidine



The compound of EXAMPLE 1 (14.55g) was dissolved in trifluoroacetic acid (100mL) and the solution stirred at room temperature under nitrogen for 1 hour. The reaction mixture was evaporated to dryness *in vacuo*. The residue was dissolved in ethyl acetate (1500mL) and washed with saturated sodium bicarbonate solution, dried over sodium sulphate and evaporated to dryness *in vacuo*. The residue was purified by column chromatography over silica gel (dichloromethane/methanol/ammonium hydroxide 85/15/2) to afford the title compound as a foam (9.77g).

TLC R_f: 0.21 (dichloromethane/methanol/ammonium hydroxide 90/10/2 by volume).

LRMS *m/z*: 515 (MH⁺).


¹H NMR : 7.32 (m, 4H), 7.20 (m, 4H), 4.62 (s, 1H), 4.43 (br s, 1H), 3.45-3.09 (m, 4H), 2.91-2.60 (m, 6H), 2.38-2.00 (m, 7H), 1.82-1.08 (m, 7H), 0.92 (m, 1H).

Found: C, 62.95; H, 6.97; N, 10.52. C₂₈H₃₆Cl₂N₄O.3/10CH₂Cl₂ requires C, 62.83; H, 6.82; N, 10.36%

[α]_D +22.0° (c=0.10, methanol).

EXAMPLE 20

3(R)-1-Benzoyl-3-(3,4-dichlorophenyl)-3-(3-[3-(homopiperazin-1-yl)azetidin-1-yl]propyl)piperidine trifluoroacetic acid salt



The compound from EXAMPLE 2 (300 mg) was dissolved in dichloromethane (7 mL), cooled in an ice/water bath and treated with trifluoroacetic acid (7 mL). After 2 h, the mixture was diluted with dichloromethane (50 mL) and 1 N sodium carbonate slowly until no effervescence was observed. The organic layer was separated (with difficulty) and washed with brine, dried over magnesium sulphate, and filtered to give a crude material (269 mg) which needed no further purification.

^1H NMR: 7.48-7.20 (m, 8H), 3.45-1.22 (m, 29H).


TLC (dichloromethane:methanol:ammonium hydroxide = 90:10:1) R_f = 0.13.

m/z 529 ($m+H$) $^+$.

The following Preparations illustrate the synthesis of certain starting materials used in the preceding Examples.

PREPARATION 1

2-(3,4-Dichlorophenyl)-5-hexenenitrile



To a solution of sodium hydride (14.8g, 60% dispersion in mineral oil) in dimethylformamide (DMF, 150ml) at 0°C under nitrogen was added a solution of 3,4-dichlorophenylacetonitrile (68.9g, 1 mol. equivalent) in DMF (300ml) and the mixture was stirred for 3 hours. A solution of 4-bromobut-1-ene (50g, 1 mol. eq.) in DMF (100ml) was then added, and the mixture was stirred at room temperature for one hour, then heated to 60°C for 5 hours.


The reaction mixture was cooled and water (1 litre) was added. The mixture was then extracted with ethyl acetate (2x500ml). The combined organic extracts were then washed with water (2 x 1 litre), dried over MgSO_4 and the solvent was removed under reduced pressure. The residue was then purified by column chromatography using silica gel, eluting with a solvent gradient of ethyl acetate : hexane (1:19 to 1:6, by volume) to give the title compound (51.5g).

TLC (hexane:ethyl acetate 6:1) R_f = 0.47.

^1H NMR : 1.85-2.1 (m,2H); 2.2-2.3 (m,2H); 3.75-3.8 (m, 1H); 5.05-5.1 (m,2H); 5.7-5.8 (m,1H); 7.15-7.2 (m, 1H); 7.4-7.45 (m,2H).

PREPARATION 2

4-Cyano-4-(3,4-dichlorophenyl)oct-7-enoic acid




To a solution of the compound of Preparation 1 (50.5g, 210.4mmol) in dioxan (150ml) at 0°C under nitrogen was added potassium tert-butoxide (1.5g, 0.06 mol.eq.) and ethyl acrylate (25.4ml, 1.11 mol.eq.), and the mixture was stirred for 1 hour. Aqueous sodium hydroxide solution (2N, 150ml) was then added and the mixture was stirred at room temperature for 70 minutes.

Methyl tert-butyl ether (300ml) was then added, and the mixture was acidified to pH 1 using aqueous 2N hydrochloric acid solution. the solution was then extracted with methyl tert-butyl ether (2 x 300 ml) and the combined organics were then dried over MgSO_4 and filtered. removal of the solvent in vacuo gave the title compound (68.12g).

^1H NMR : 1.8-2.6 (m,9H), 4.9-5.0 (m,2H), 5.65-5.75 (M,1H), 7.2-7.25 (m,1H), 7.45-7.5 (m,2H).

PREPARATION 3

(+)-4S-4-Cyano-4-(3,4-dichlorophenyl)oct-7-enoic acid



Part a)

A mixture of the compound of PREPARATION 2 (247g) in acetone (247ml) and S-(-)-1-(1-naphthyl)ethylamine (68.8g) was heated to 50°C and warm hexane (1200ml) added until cloudiness appeared. The mixture was warmed back to 50°C and sufficient additional hexane (50ml) added to induce cloudiness again. The resulting cloudy solution was stirred at room temperature for 18 hours. The sticky precipitate was collected by filtration and washed with hexane (3x200ml) and acetone/hexane (200ml; 10% by volume). The solid was dried at 50°C to afford the 1-naphthyl)ethylamine salt of the title compound, 108.7g. This material was stirred in acetone (300ml) and the small amount of insoluble material removed by filtration. The filtrate was stirred at room temperature and hexane (1200ml) added to the cloud point. Precipitation occurred within 5 minutes and, after stirring at room temperature for 1 hour, the solid precipitate was collected by filtration and dried (54.1g). The process was repeated with 210.3g of the compound of PREPARATION 2 and (S)-(-)-1-(1-naphthyl)ethylamine (58.7g) to afford a solid which was combined with the 54.1g of product obtained in the first run to give a total yield of 135.2g.

Part b)


The salt from part a) was stirred in dichloromethane (1000ml) and water (1000ml) and basified with 40% sodium hydroxide solution. The mixture was stirred for 30 minutes to afford a stable emulsion which was treated with solid sodium chloride. After a further period of stirring (15 minutes) the layers were separated and the aqueous solution was diluted with water (1000ml) and washed with further dichloromethane (1000ml). The combined organic washes were washed with saturated brine solution and evaporated to dryness to recover the chiral resolving agent (45g). The aqueous phase was stirred with dichloromethane (1500ml) and acidified to pH <0.5 with concentrated hydrochloric acid and the resulting mixture was stirred for a further 30 minutes. The layers were separated and the aqueous layer extracted with dichloromethane (500ml). The combined organic extracts were washed with water, dried over sodium sulphate and evaporated to dryness *in vacuo* to afford the title compound as a white solid, (69.8g).

TLC R_f: 0.19 (dichloromethane/methanol/ammonium hydroxide 85/15/2 by volume).

m/z: 329 (MNH₄⁺).

¹H NMR: 7.43 (m, 2H), 7.21 (m, 1H), 5.70 (m, 1H), 4.97 (m, 2H), 2.60-1.80 (m, 9H).

[α]_D -11.4° (c=0.45, methanol)

PREPARATION 4**(-)-4R-4-Cyano-4-(3,4-dichlorophenyl)-7-hydroxyheptanoic acid**

Ozone in a stream of oxygen was bubbled through a solution of the compound of Preparation 3 (15.6g) in ethanol (400mL) at -78°C for 1 hour. The cooling was removed and the resulting blue/green solution was purged with nitrogen for 1 hour to remove excess ozone. The solution was then cooled in an ice-bath and sodium borohydride (9.45g) was added portionwise and the resulting mixture was stirred at room temperature for 18 hours. The solution was cooled in an ice-bath and the solution acidified to pH 2 by the cautious addition of 5N hydrochloric acid. The partially worked up reaction was combined with the partially worked up material from an identical preparation and diluted with water (300mL). The resulting solution was extracted into dichloromethane (3x1000mL). The combined organic extracts were dried over magnesium sulphate and evaporated *in vacuo* to afford the title compound as a foam (30.0g) which was used without any further purification.


TLC R_f : 0.11 (dichloromethane/methanol/ammonium hydroxide 80/20/3 by volume).

m/z : 333 (MNH_4^+).

^1H NMR: 7.48 (m, 2H), 7.24 (m, 1H), 4.40 (broad s, 2H), 3.62 (t, 2H), 2.62-1.97 (m, 6H), 1.72 (m, 1H), 1.38 (m, 1H).

$[\alpha]_D -10.2^{\circ}$ ($c=0.10$, methanol).

PREPARATION 5**(+)-3(R)-3-(3-Acetyloxypropyl)-3-(3,4-dichlorophenyl)glutarimide**



Concentrated sulphuric acid (3mL) was added to a solution of the compound of PREPARATION 4 (29.7g) in glacial acetic acid (400mL), and the mixture heated under reflux for 6 hours. The reaction mixture was cooled to room temperature and allowed to stand for 18 hours. Evaporation of the volatiles *in vacuo* afforded a residue which was dissolved in dichloromethane (1000mL) and washed with saturated sodium hydrogen carbonate solution and saturated brine. The organic solution was dried over sodium sulphate and evaporated to dryness *in vacuo* to afford the title compound as an off white solid, (31.48g) which was used without further purification.

TLC R_f: 0.55 (dichloromethane/methanol 95/5 by volume).


m/z: 375 (MNH₄⁺)

¹H NMR : 7.85 (broad s, 1H), 7.44 (d, 1H), 7.34 (s, 1H), 7.09 (d, 1H), 3.98 (t, 2H), 2.62 (m, 1H), 2.34 (m, 2H), 2.21 (m, 1H), 1.95 (m, 5H), 1.57 (m, 2H).

[α]_D +108.2° (c=0.118, methanol).

PREPARATION 6

(-)-3(S)-3-(3,4-Dichlorophenyl)-3-(3-hydroxypropyl)piperidine



Borane dimethylsulphide complex (83mL) was added dropwise over 10 minutes to a solution of the compound of PREPARATION 5 (31.27g) in dry tetrahydrofuran (450mL) at 0°C. The mixture was allowed to warm to room temperature over 30 minutes and then to reflux over an hour. After 2 hours at reflux the solution was cooled in an ice-bath and methanolic hydrogen chloride (300mL of an approximately 1.75N solution) was added dropwise with caution over a period of 1 hour. The strongly acidic solution was allowed to warm to room temperature and

stirred at this temperature overnight. The solution was evaporated to dryness *in vacuo* and the residue was partitioned between water (500mL) and diethyl ether (500mL). The aqueous solution was washed with further diethyl ether (500mL) and the combined organics discarded. The aqueous solution was basified with ammonium hydroxide (s.g. 0.880) and extracted into diethyl ether (2x750mL). The combined extracts were dried over sodium sulphate and evaporated to dryness to afford the title compound as a gum, 25.2g which was used without further purification. A small quantity (650mg) was purified by column chromatography over silica gel (dichloromethane/methanol/ammonium hydroxide) to afford a pure sample of the title compound, (546mg).

TLC R_f: 0.34 (dichloromethane/methanol/ammonium hydroxide 90/10/2 by volume).


LRMS *m/z*: 288 (MH⁺).

¹H NMR : 7.37 (m, 2H), 7.12 (d, 1H), 3.45 (t, 2H), 3.18 (d, 1H), 2.80 (m, 3H), 2.01 (m, 1H), 1.70 (m, 1H), 1.59 (m, 5H), 1.42 (m, 1H), 1.15 (m, 2H).

[α]_D -3.7° (c=0.107, methanol)

PREPARATION 7

(+)-3(S)-1-Benzoyl-3-(3,4-dichlorophenyl)-3-(3-hydroxypropyl)piperidine



To an ice-cold solution of the compound of PREPARATION 6 (15.44g) and triethylamine (13.01g) in dry dichloromethane (300mL) was added benzoyl chloride (16.57g) slowly over about 10 minutes. The resulting mixture was allowed to warm to room temperature and was stirred for 24 hours. The reaction mixture was evaporated to dryness *in vacuo* and the residue was dissolved in ethanol (300mL) and dichloromethane (100mL). To this solution was added sodium hydroxide (155mL of 2N aqueous solution) and the resulting mixture was stirred at room temperature for 5 hours. The solution was cooled in an ice-bath and acidified to pH6 using 6N hydrochloric acid and then evaporated to low volume. The mixture was extracted with ethyl acetate (2x300mL) and the combined extracted dried over sodium sulphate and evaporated to dryness *in vacuo*. The residue was purified by column chromatography over

silica gel (dichloromethane/methanol/ammonium hydroxide 95/5/0.5 by volume) to afford the title compound as a white foam (19.90g).


TLC R_f: 0.32 (dichloromethane/methanol/ammonium hydroxide 95/5/0.5 by volume)

m/z: 393 (MH⁺)

[α]_D +35.1° (c=0.13, methanol)

PREPARATION 8

(+)-3(S)-1-Benzoyl-3-(3,4-dichlorophenyl)-3-(3-methanesulphonyloxypropyl)piperidine



Methanesulphonyl chloride (4.63mL) was added over a period of 10 minutes to an ice-cold solution of the compound of PREPARATION 7 (19.57g) and triethylamine (9.74mL) under nitrogen. The reaction mixture was stirred at 0°C for 30 minutes and then warmed to room temperature. The reaction was diluted with dichloromethane (150mL) and washed successively with water, 1N aqueous citric acid solution and saturated aqueous sodium bicarbonate solution. The organic phase was dried over sodium sulphate and evaporated to dryness *in vacuo* to afford the title compound as a foam (23.06g).

TLC R_f: 0.61 silica, dichloromethane/methanol 95/5 by volume.


LRMS *m/z*: 471 (MH⁺).

¹H NMR : 7.39 (m, 5H), 7.24 (m, 3H), 4.35 (m, 1H), 4.09 (m, 2H), 3.58 (d, 1H), 3.40 (m, 2H), 2.95 (s, 3H), 2.12 (m, 1H), 1.93-1.28 (m, 7H).

[α]_D +34.2° (c=0.168, methanol)

PREPARATION 9

3-(4-tert-Butoxycarbonylpiperazin-1-yl)-1-diphenylmethylazetidine




A mixture of N-*tert*-butoxycarbonylpiperazine (14.08g), potassium carbonate (26.1g), and 1-diphenylmethyl-3-methanesulphonyloxyazetidine (WO 96/05193)(20g) in dry acetonitrile (600mL) was heated under reflux for 32 hours. The reaction mixture was cooled to room temperature, diluted with ethyl acetate and washed with water and brine. The organic phase was dried over sodium sulphate and evaporated to dryness *in vacuo*. The residue was purified by column chromatography over silica (dichloromethane/methanol 97/3 by volume) to afford the title compound as a solid (22.53g).

TLC R_f: 0.17 (dichloromethane/methanol 98/2 by volume).

LRMS *m/z*: 408 (MH⁺).

PREPARATION 10

3-(4-*tert*-Butoxycarbonylpiperazin-1-yl)azetidine




1-Chloroethyl chloroformate (6.88mL) was added dropwise to an ice-cold solution of the compound of PREPARATION 9 (22g) in dry dichloromethane (100mL). The resulting mixture was stirred at room temperature for 3 hours, before the volatiles were evaporated *in vacuo*. To the residue was added powdered potassium carbonate (13.2g) and methanol (125mL) and the mixture was heated under reflux for 2 hours. The cooled reaction mixture was filtered through SOLKAFLOCTM filter aid and the filtrate was evaporated to dryness *in vacuo*. The residue was purified by column chromatography over silica gel (dichloromethane/methanol/ammonium hydroxide 80/20/3) to afford the title compound (14g). TLC R_f: 0.14 (dichloromethane/methanol/ammonium hydroxide 90/10/2 by volume).

LRMS *m/z*: 242 (MH⁺).

PREPARATION 11

1-Diphenylmethyl-3-(4-*tert*-butoxycarbonylhomopiperazin-1-yl)azetidine




To a solution of 1-diphenylmethyl-3-methanesulphonyloxyazetidine (WO 96/05193) (25 g) in acetonitrile (800 mL) were added the compound of PREPARATION 24 (19 g) and potassium carbonate (33 g). The reaction mixture was heated to reflux for 12 h, then cooled to room temperature. Water was added to the resulting slurry, and the mixture extracted three times with ethyl acetate. The combined organics were washed with brine, dried over magnesium sulphate, filtered and dried *in vacuo* to yield crude product. Chromatography on silica gel eluting with dichloromethane:methanol (95:5) yielded the title compound (24.7 g).

^1H NMR : 7.41-7.14 (m, 10H), 4.39 (s, 1H), 3.41 (m, 6H), 3.15 (m, 1H), 2.82 (m, 2H), 2.38 (m, 4H), 1.78 (m, 2H), 1.45 (s, 9H).

LRMS m/z 422 ($m+H$) $^+$.

PREPARATION 12

3-(4-*tert*-Butoxycarbonylhomopiperazin-1-yl)azetidine




The compound from PREPARATION 11 (1.0 g) was dissolved in ethanol (75 mL) and mixed with palladium hydroxide (Pearlman's catalyst, 200 mg), then subjected to hydrogen gas at 345 kPa (50 p.s.i.) for 16 h. The catalyst was removed by filtration and the filtrate evaporated to dryness. Chromatography on silica gel with dichloromethane:methanol:ammonium hydroxide (90:10:2) as eluant yielded the title product.

^1H NMR : 3.55-3.38 (m, 10H), 2.40 (m, 4H), 1.82 (m, 2H), 1.46 (s, 9H).

m/z 256 ($m+H$) $^+$.

PREPARATION 13

(+)-3(R)-1-Benzoyl-3-(3,4-dichlorophenyl)-3-(3-[3-piperazinoazetidin-1-yl]propyl)piperidine



The compound of EXAMPLE 1 (14.55g) was dissolved in trifluoroacetic acid (100mL) and the solution stirred at room temperature under nitrogen for 1 hour. The reaction mixture was evaporated to dryness *in vacuo*. The residue was dissolved in ethyl acetate (1500mL) and washed with saturated sodium bicarbonate solution, dried over sodium sulphate and evaporated to dryness *in vacuo*. The residue was purified by column chromatography over silica gel (dichloromethane/methanol/ammonium hydroxide 85/15/2) to afford the title compound as a foam, 9.77g.

TLC R_f: 0.21 (dichloromethane/methanol/ammonium hydroxide 90/10/2 by volume).

LRMS *m/z*: 515 (MH⁺).


¹H NMR : 7.32 (m, 4H), 7.20 (m, 4H), 4.62 (s, 1H), 4.43 (br s, 1H), 3.45-3.09 (m, 4H), 2.91-2.60 (m, 6H), 2.38-2.00 (m, 7H), 1.82-1.08 (m, 7H), 0.92 (m, 1H).

Found: C, 62.95; H, 6.97; N, 10.52. C₂₈H₃₆Cl₂N₄O.3/10CH₂Cl₂ requires C, 62.83; H, 6.82; N, 10.36%

[α]_D +22.0° (c=0.10, methanol).

PREPARATION 14

3(R)-1-Benzoyl-3-(3,4-dichlorophenyl)-3-(3-[3-(homopiperazin-1-yl)azetidin-1-yl]propyl)piperidine trifluoroacetic acid salt



The compound from EXAMPLE 2 (300 mg) was dissolved in dichloromethane (7 mL), cooled in an ice/water bath and treated with trifluoroacetic acid (7 mL). After 2 h, the mixture was diluted with dichloromethane (50 mL) and 1 N sodium carbonate slowly until no effervescence was observed. The organic layer was separated (with difficulty) and washed with brine, dried

over magnesium sulphate, and filtered to give a crude material (269 mg) which needed no further purification.


^1H NMR: 7.48-7.20 (m, 8H), 3.45-1.22 (m, 29H).

TLC (dichloromethane:methanol:ammonium hydroxide = 90:10:1) R_f = 0.13.

LRMS m/z 529 (m+H) $^+$.

PREPARATION 15

3(S)-1-*tert*-Butoxycarbonyl-3-(3,4-dichlorophenyl)-3-(3-hydroxypropyl)piperidine




The compound from PREPARATION 6 (5.0 g) and *tert*-butoxycarbonyl anhydride were dissolved in dichloromethane (50 mL) and stirred at ambient temperature overnight. The solution was washed with aqueous sodium bicarbonate and brine, dried over magnesium sulphate, filtered, and evacuated to dryness to give the product (5.59 g).

^1H NMR : 7.43 (s, 1H), 7.39 (d, J = 8Hz, 1H), 7.18 (d, J = 8Hz, 1H), 3.98 (m, 1H), 3.78-3.02 (m, 6H), 2.03 (m, 1H), 1.80-1.03 (m, 17H).

m/z 389 (m+H) $^+$.

PREPARATION 16

3(S)-1-*tert*-Butoxycarbonyl-3-(3,4-dichlorophenyl)-3-(3-methanesulphonyloxypropyl)piperidine



Methanesulphonyl chloride (1.22 mL) was added dropwise to a solution of the compound from PREPARATION 15 (5.58 g) and triethylamine (3.0 mL) in dichloromethane (50 mL) while cooling in an ice bath. The mixture was then allowed to warm to ambient temperature for 1.5 hours before quenching with aqueous sodium bicarbonate. The extracted organic layer


was washed with brine, dried over magnesium sulphate, filtered, and evacuated to dryness to give the product (6.37 g).

^1H NMR : 7.41 (m, 2H), 7.19 (m, 1H), 4.04 (t, $J = 7\text{Hz}$, 2H), 4.02 (m, 1H), 3.70-3.10 (m, 3H), 2.93 (s, 3H), 2.03 (m, 1H), 1.80-1.22 (m, 12H).

LRMS m/z 270 ($[\text{m} - \text{HOSO}_2\text{Me}] + \text{H}$) $^+$.

PREPARATION 17

3(R)-1-*tert*-Butoxycarbonyl-3-(3,4-dichlorophenyl)-3-(3-[3-(4-methanesulphonylpiperazin-1-yl)azetidin-1-yl]propyl)piperidine.




The compound from PREPARATION 16 (1.0g) was mixed with the compound from PREPARATION 22 (603 mg) and potassium hydrogen carbonate (643 mg) in acetonitrile (30 mL) and the mixture was heated to reflux overnight. The reaction mixture was cooled to ambient temperature and partitioned between ethyl acetate (50 mL) and water (25 mL). The organic layer was then washed with brine, dried over magnesium sulphate, filtered, and evaporated *in vacuo* to yield a crude mixture. Purification on silica gel with dichloromethane:methanol:ammonium hydroxide (98:2:1) yielded the desired compound (1.18 g).

^1H NMR : 7.42-7.34 (m, 2H), 7.12 (m, 1H), 4.02 (m, 1H), 3.71-1.90 (m, 22H), 1.75-0.78 (m, 17H).

m/z 591 ($\text{m} + \text{H}$) $^+$.

PREPARATION 18

3(S)-3-(3,4-dichlorophenyl)-3-(3-[3-(4-methanesulphonylpiperazin-1-yl)azetidin-1-yl]propyl)piperidine




The compound from PREPARATION 17 (606 mg) was dissolved in dichloromethane (2 mL) and cooled in an ice water bath before trifluoroacetic acid (2 mL) was added. After 1 h, toluene (10 mL) was added followed by solvent evaporation *in vacuo*. To the residue was added 0.2 N sodium hydroxide (50 mL), and the mixture was extracted repeatedly with dichloromethane. The combined organic phases were dried over magnesium sulphate, filtered, and evaporated to give a white solid (376 mg).

^1H NMR : 7.38 (m, 2H), 7.12 (m, 1H), 3.31-1.82 (m, 22H), 1.8-1.38 (m, 6H), 0.93 (m, 2H).
 m/z 491 ($m+H$) $^+$.

PREPARATION 19


1-Diphenylmethyl-3-(N-methyl benzenesulphonamido)azetidine



To a solution of 1-diphenylmethyl-3-methanesulphonyloxyazetidine (WO 96/05193)(1.4 g) in dioxane (25 mL) were added N-methyl benzenesulphonamide (1.2 g) and caesium carbonate (2.6 g). The reaction mixture was heated to reflux for 3 h, then cooled to room temperature. Water (20 mL) was added to the resulting slurry, and the mixture was extracted with ethyl acetate (3 x 25 mL). The combined organics were washed with brine, dried over magnesium sulphate, filtered and dried *in vacuo* to yield crude product. Chromatography on silica gel eluting with ethyl acetate:hexanes (2:3 by volume) yielded the title compound as a white foam (1.31 g).


^1H NMR : 7.68 (d, $J = 8$ Hz, 2H), 7.54 (t, $J = 8$ Hz, 1H), 7.46 (t, $J = 8$ Hz, 2H), 7.32 (d, $J = 7$ Hz, 4H), 7.22 (t, $J = 7$ Hz, 4H), 7.14 (t, $J = 7$ Hz, 2H), 4.27 (s, 1H), 3.92 (quintet, $J = 7$ Hz, 1H), 3.33 (t, $J = 7$ Hz, 2H), 2.99 (t, $J = 7$ Hz, 2H), 2.66 (s, 3H).

m/z 393 ($m+H$) $^+$

PREPARATION 20**3-(N-Methyl benzenesulphonamido)azetidine**

The compound from PREPARATION 19 (1.3 g) was dissolved in ethanol (30 mL) and mixed with palladium hydroxide (Pearlman's catalyst, 600 mg), then subjected to hydrogen gas at 345 kPa (50 p.s.i.) for 24 h. The catalyst was removed by filtration and the filtrate evaporated to dryness. Chromatography on silica gel with dichloromethane:methanol:ammonium hydroxide (93:7:1) as eluant yielded the title compound.

^1H NMR : 7.70 (d, $J = 7\text{Hz}$, 2H), 7.61 (t, $J = 7\text{Hz}$, 1H), 7.53 (t, $J = 7\text{Hz}$, 2H), 4.60 (m, 1H), 4.06 (dd, $J = 11$ and 8Hz , 2H), 3.99 (dd, $J = 11$ and 8Hz , 2H), 3.45 (s, 1H), 2.76 (s, 3H).
 m/z 227 ($m+H$) $^+$.


PREPARATION 21**1-Diphenylmethyl-3-(N-phenyl methanesulphonamido)azetidine**

To a solution 1-diphenylmethyl-3-methanesulphonyloxyazetidine (WO 96/05193) (1.2 g) in dioxane (15 mL) was added N-phenyl methanesulphonamide (1.0 g, 5.85 mmol) and caesium carbonate (2.0 g, 6.1 mmol). The reaction mixture was heated to reflux for 2 h, then cooled to room temperature. Water (20 mL) was added to the resulting slurry, and the mixture was extracted with magnesium sulphate, filtered and dried *in vacuo* to yield crude product.

Rescrystallization from ethyl acetate:hexanes (4:1) gave 964 mg of a white solid.

^1H NMR : 7.38-7.09 (m, 15H), 4.55 (m, 1H), 4.17 (s, 1H), 3.47 (dd, $J = 7$ and 3Hz , 2H), 2.81 (dd, $J = 7$ and 3Hz , 2H), 2.76 (s, 3H).

m/z 393 ($m+H$) $^+$.


PREPARATION 22**3-(N-Phenyl methanesulphonamido)azetidine**

The compound from PREPARATION 21 (890 mg) was dissolved in ethanol (30 mL) and mixed with palladium hydroxide (Pearlman's catalyst, 800 mg), then subjected to hydrogen gas at 345 kPa (50 p.s.i.) for 48 h. The catalyst was removed by filtration and the filtrate was evaporated to dryness. Chromatography on silica gel with

dichloromethane:methanol:ammonium hydroxide (93:7:1) as eluant yielded pure product.

^1H NMR : 7.41-7.30 (m, 3H), 7.21 (m, 2H), 4.84 (m, 1H), 3.59 (m, 4H), 2.80 (s, 3H), 1.69 (br s, 1H)

m/z 227 ($m+H$)⁺


PREPARATION 22**1-Diphenylmethyl-3-(4-methanesulphonylpiperazinyl)azetidine**

To a solution 1-diphenylmethyl-3-methanesulphonyloxyazetidine (WO 96/05193) (7.73 g) in acetonitrile (250 mL) was added methanesulphonylpiperazine (4.8 g). The reaction mixture was heated to reflux overnight, then cooled to room temperature and concentrated to a foam. Purification on silica gel with dichloromethane:methanol:ammonium hydroxide (95:5:0.5) as eluant yielded pure product (5.97 g).

TLC (dichloromethane:methanol:ammonium hydroxide = 90:10:1) R_f = 0.44

m/z 386 ($m+H$)⁺

PREPARATION 23**3-(4-Methanesulphonylpiperazine)azetidine**




The compound from PREPARATION 22 (2.77 g) was dissolved in anhydrous dichloromethane (40 mL), cooled in an ice bath, and treated with 1-chloroethyl chloroformate (2.06 g). The solution was allowed to warm to ambient temperature and was stirred for 3 h before removing the solvent *in vacuo*. To the residue was added methanol (40 mL) and the solution was heated to reflux for 1 h. After cooling to ambient temperature and stirring overnight, the resulting precipitate was collected by filtration.

$^1\text{H NMR}$: 4.32 (m, 2H), 4.11 (m, 1H), 3.99 (m, 2H), 3.38 (m, 4H), 2.95 (s, 3H), 2.21 (m, 4H)
 m/z 221 ($m+H$)⁺

PREPARATION 24

4-tert-Butoxycarbonylhomopiperazine



tert-Butoxycarbonyl anhydride (195 g) was dissolved in dichloromethane (350 mL) and added dropwise to a solution of homopiperazine (100 g) and triethylamine (210 mg) in dichloromethane (500 mL) while cooling in an ice/water bath. The mixture was stirred overnight, allowing to warm to ambient temperature. After removing the solvent *in vacuo*, the crude mixture was partitioned between 2 N citric acid (aqueous) and diethyl ether. The aqueous layer was washed with diethyl ether, then basified with 2 N sodium hydroxide and extracted repeatedly with dichloromethane. The combined dichloromethane layers were combined, dried over magnesium sulphate, filtered, and evaporated to dryness to yield a yellow oil (170 g).

TLC (dichloromethane:methanol:ammonium hydroxide = 90:10:1) R_f = 0.43

m/z 201 ($m+H$)⁺.

PREPARATION 25

3-morpholinoazetidine dihydrochloride

A mixture of 1-diphenylmethyl-3-morpholinoazetidine (prepared in an analogous manner to the compound of preparation 22 above) (18.6g), palladium hydroxide (2g), ethanol (200ml)

and 1N aqueous hydrochloric acid solution (52ml) was stirred under an atmosphere of hydrogen at 345 kPa (50p.s.i.) for 3 days. The catalyst was then removed by filtration and the filtrate evaporated to dryness. Addition of dichloromethane (100ml) to the residue and trituration gave a solid which was recrystallised from methanol to give the title compound (10.2g).

m/z 179 ($m+H$)⁺.


PHARMACOLOGICAL DATA

The data below illustrates the *in vitro* affinity of substances of the invention for the guinea pig cortex NK₃ receptor, which assays were carried out according to the method mentioned earlier on page 15.

<u>Example</u>	<u>pIC₅₀</u>
3	8.8
13	7.95
17	8.4

CLAIMS

1. A compound of the formula (I):



and pharmaceutically-acceptable salts thereof,

wherein

A is CO, SO₂ or CH₂;

Ar¹ is phenyl optionally substituted by one or more substituents independently selected from halogen, C₁₋₆ alkoxy optionally substituted by one or more halogen, and C₁₋₆ alkyl optionally substituted by one or more halogen;


X¹ is C₃₋₇ cycloalkyl, aryl or C₁₋₆ alkyl,

said C₁₋₆ alkyl being optionally substituted by fluoro, CO₂H, CO₂(C₁₋₄alkyl), C₃₋₇ cycloalkyl, adamantyl, aryl or het,


and said C₃₋₇ cycloalkyl being optionally substituted by 1 or 2 substituents each independently selected from C₁₋₄ alkyl, C₃₋₇ cycloalkyl, C₁₋₄ alkoxy, OH, F, fluoro(C₁₋₄ alkyl) and fluoro(C₁₋₄ alkoxy);

X is a direct link or NR¹;

R is SO₂aryl, SO₂(C₁₋₆ alkyl optionally substituted by one or more halogen),




or




R^1 is H, C_{1-6} alkyl optionally substituted by one or more halogen,
or R^1 is phenyl optionally substituted by one or more substituents independently selected from halogen, C_{1-6} alkoxy optionally substituted by one or more halogen, or C_{1-6} alkyl optionally substituted by one or more halogen;

Y is O, NCO(C_{1-6} alkyl optionally substituted by one or more halogen), NCO₂(C_{1-6} alkyl optionally substituted by one or more halogen), NSO₂(C_{1-6} alkyl optionally substituted by one or more halogen), NCOaryl, NCO₂aryl, NSO₂aryl, NSO₂(C_{1-6} alkyl optionally substituted by one or more halogen), CH₂, CHF, CF₂, NH, NCH₂aryl, N(C_{1-6} alkyl optionally substituted by one or more halogen), or NCH₂(C_{3-7} cycloalkyl);

with the proviso that X is not NR¹ when R is




or




wherein "aryl" used in all the above definitions represents phenyl or naphthyl optionally substituted with one or more substituents independently selected from halogen, C_{1-6} alkoxy optionally substituted by one or more halogen, or C_{1-6} alkyl optionally substituted by one or more halogen;

and "het" used in the definition of X¹ means thienyl or a 5- or 6-membered ring heteroaryl group containing either 1 or 2 nitrogen heteroatoms or 1 nitrogen heteroatom and one oxygen or sulphur heteroatom, each optionally substituted by 1 or 2 substituents each independently selected from C_{1-4} alkyl, C_{1-4} alkoxy, halogen, fluoro(C_{1-4} alkyl) and fluoro(C_{1-4} alkoxy).

2. A compound or salt according to claim 1 wherein A is CO or SO₂.
3. A compound or salt according to any preceding claim wherein Ar¹ is phenyl optionally substituted by one or more halogen atoms.
4. A compound or salt according to any preceding claim wherein X¹ is aryl or C₁₋₆ alkyl, said C₁₋₆ alkyl being optionally substituted by fluoro, CO₂H, CO₂(C₁₋₄)alkyl, C₃₋₇ cycloalkyl, adamantyl, aryl or het, and said C₃₋₇ cycloalkyl being optionally substituted by 1 or 2 substituents each independently selected from C₁₋₄ alkyl, C₃₋₇ cycloalkyl, C₁₋₄ alkoxy, OH, F, fluoro(C₁₋₄ alkyl) and fluoro(C₁₋₄ alkoxy).
5. A compound or salt according to any preceding claim wherein X is a direct link or NR¹, where R¹ is C₁₋₆ alkyl optionally substituted by one or more halogen, or R¹ is phenyl optionally substituted by one or more halogen substituents.
6. A compound or salt according to any preceding claim wherein R is SO₂aryl, SO₂(C₁₋₆ alkyl optionally substituted by one or more halogen),




or




where Y is O, NCO(C₁₋₆ alkyl optionally substituted by one or more halogen), NCO₂(C₁₋₆ alkyl optionally substituted by one or more halogen), NSO₂(C₁₋₆ alkyl optionally substituted by one or more halogen), NCOaryl, CH₂, CHF, CF₂, NH, NCH₂aryl, N(C₁₋₆ alkyl optionally substituted by one or more halogen), or NCH₂(C₃₋₇ cycloalkyl).

7. A compound or salt according to any preceding claim wherein A is CO.

8. A compound or salt according to any preceding claim wherein Ar^1 is phenyl optionally substituted by up to two halogen atoms.
9. A compound or salt according to any preceding claim wherein X^1 is C_{1-6} alkyl substituted by C_{3-7} cycloalkyl, said C_{3-7} cycloalkyl being optionally substituted by 1 or 2 substituents each independently selected from C_{1-4} alkyl, C_{3-7} cycloalkyl, C_{1-4} alkoxy, OH, F, fluoro(C_{1-4} alkyl) and fluoro(C_{1-4} alkoxy), or X^1 is phenyl optionally substituted by one or more substituents independently selected from halogen and C_{1-6} alkyl optionally substituted by one or more halogen.
10. A compound or salt according to any preceding claim wherein X is a direct link, N-methyl or N-phenyl.
11. A compound or salt according to any preceding claim wherein R is SO_2 (phenyl optionally substituted with one or more substituents independently selected from halogen, C_{1-6} alkoxy optionally substituted by one or more halogen, or C_{1-6} alkyl optionally substituted by one or more halogen), SO_2 (C_{1-3} alkyl optionally substituted by one or more halogen),



or




where Y is O, $\text{NCO}(\text{C}_{1-6} \text{ alkyl})$, $\text{NCO}_2(\text{C}_{1-6} \text{ alkyl})$, $\text{NSO}_2(\text{C}_{1-6} \text{ alkyl}$ optionally substituted by one or more halogen), NCO (phenyl optionally substituted with one or more substituents independently selected from halogen, C_{1-6} alkoxy optionally substituted by one or more halogen, or C_{1-6} alkyl optionally substituted by one or more halogen), CHF, CF_2 , NH, NCH_2 (phenyl optionally substituted with one or more substituents independently selected from halogen, C_{1-6} alkoxy optionally substituted by one or more halogen, or C_{1-6} alkyl optionally substituted by one or more halogen), $\text{N}(\text{C}_{1-6} \text{ alkyl})$, or $\text{NCH}_2(\text{C}_{3-5} \text{ cycloalkyl})$.


12. A compound or salt according to any preceding claim wherein Ar^1 is phenyl optionally substituted by one or two chlorine atoms.

13. A compound or salt according to any preceding claim wherein X^1 is C_{1-6} alkyl substituted by C_{3-7} cycloalkyl, or is phenyl optionally substituted by one or more substituents independently selected from halogen and C_{1-3} alkyl optionally substituted by one or more halogen.

14. A compound or salt according to any preceding claim wherein R is SO_2Ph , SO_2CH_3 ,



or




where Y is O, NCOCH_3 , $\text{NCO}_2\text{C}(\text{CH}_3)_3$, NSO_2CH_3 , $\text{NSO}_2\text{C}_2\text{H}_5$, NSO_2Ph , $\text{NSO}_2\text{CH}_2\text{Ph}$, $\text{NSO}_2\text{CH}(\text{CH}_3)_2$, $\text{NSO}_2\text{CH}_2\text{CF}_3$, NCOPh , CHF , NH , NCH_2Ph , or N-cyclopropylmethyl.


15. A compound or salt according to any preceding claim wherein Ar^1 is phenyl, 4-chlorophenyl or 3,4-dichlorophenyl.

16. A compound or salt according to any preceding claim wherein X^1 is (C_{3-7} cycloalkyl)methyl, or phenyl optionally substituted by one or more substituents independently selected from halogen and methyl optionally substituted by one or more halogen.

17. A compound or salt according to any preceding claim wherein R is SO_2Ph , SO_2CH_3 ,



or



where Y is O, NCOCH₃, NCO₂C(CH₃)₃, NSO₂CH₃, NSO₂C₂H₅, NSO₂Ph, NSO₂CH₂Ph, NSO₂CH(CH₃)₂, NSO₂CH₂CF₃, NCOPh, or CHF.

18. A compound or salt according to any preceding claim wherein X¹ is cyclopropylmethyl or phenyl optionally substituted by one or more fluorine or chlorine atoms.

19. A compound or salt according to any preceding claim wherein X¹ is cyclopropylmethyl, phenyl, 3-chlorophenyl or 3,4-difluorophenyl.

20. A compound or salt according to claim 1 which is selected from:

- (+)-3(R)-1-benzoyl-3-(3-[3-(4-*tert*-butoxycarbonylpiperazin-1-yl)azetidin-1-yl]propyl)-3-(3,4-dichlorophenyl)piperidine;
- 3(R)-1-benzoyl 3-(3-[3-(4-*tert*-butoxycarbonylhomopiperazin-1-yl)azetidin-1-yl]propyl)-3-(3,4-dichlorophenyl)piperidine;
- (+)-3(R)-1-benzoyl-3-(3,4-dichlorophenyl)-3-(3-[3-(4-methanesulphonylpiperazin-1-yl)azetidin-1-yl]propyl)piperidine;
- (+)-3(R)-3-(3-[3-(4-acetylpiperazin-1-yl)azetidin-1-yl]propyl)-1-benzoyl-3-(3,4-dichlorophenyl)piperidine;
- (+)-3(R)-1-benzoyl-3-(3-[3-(4-benzoylpiperazin-1-yl)azetidin-1-yl]propyl)-3-(3,4-dichlorophenyl)piperidine;
- (+)-3(R)-1-benzoyl-3-(3,4-dichlorophenyl)-3-(3-[3-(4-methanesulphonylpiperazin-1-yl)azetidin-1-yl]propyl)piperidine;
- (+)-3(R)-3-(3-[3-(4-benzenesulphonylpiperazin-1-yl)azetidin-1-yl]propyl)-1-benzoyl-3-(3,4-dichlorophenyl)piperidine;
- (+)-3(R)-1-benzoyl-3-(3-[3-(4-benzylsulphonylpiperazin-1-yl)azetidin-1-yl]propyl)-3-(3,4-dichlorophenyl)piperidine;
- (+)-3(R)-1-benzoyl-3-(3,4-dichlorophenyl)-3-(3-[3-(4-*iso*-propylsulphonylpiperazin-1-yl)azetidin-1-yl]propyl)piperidine;
- (+)-3(R)-1-benzoyl-3-(3,4-dichlorophenyl)-3-(3-[3-(4-(2,2,2-trifluoroethane)sulphonylpiperazin-1-yl)azetidin-1-yl]propyl)piperidine;
- (+)-3(R)-1-benzoyl-3-(3,4-dichlorophenyl)-3-(3-[3-morpholinoazetidin-1-yl]propyl)piperidine;
- (+)-3(R)-1-benzoyl-3-(3,4-dichlorophenyl)-3-(3-[3-(N-methyl benzenesulphonamido)azetidin-1-yl]propyl)piperidine;

(+)-3(R)-1-benzoyl-3-(3,4-dichlorophenyl)-3-(3-[3-(N-phenyl methanesulphonamido)azetidin-1-yl]propyl)piperidine;
3(R)-1-benzoyl-3-(3,4-dichlorophenyl)-3-(3-[3-(4-methanesulphonylhomopiperazin-1-yl)azetidin-1-yl]propyl)piperidine;
3(R)-1-benzenesulphonyl-3-(3,4-dichlorophenyl)-3-(3-[3-(4-methanesulphonylpiperazin-1-yl)azetidin-1-yl]propyl)piperidine;
(+)-3(R)-1-(cyclopropylacetyl)-3-(3,4-dichlorophenyl)-3-(3-[3-(4-methanesulphonylpiperazin-1-yl)azetidin-1-yl]propyl)piperidine;
(+)-3(R)-3-(3,4-dichlorophenyl)-1-(3,4-difluorobenzoyl)-3-(3-[3-(4-methanesulphonylpiperazin-1-yl)azetidin-1-yl]propyl)piperidine;
(+)-3(R)-1-(3-chlorobenzoyl)-3-(3,4-dichlorophenyl)-3-(3-[3-(4-methanesulphonylpiperazin-1-yl)azetidin-1-yl]propyl)piperidine;
(+)-3(R)-1-benzoyl-3-(3,4-dichlorophenyl)-3-(3-[3-piperazinoazetidin-1-yl]propyl)piperidine;
and 3(R)-1-benzoyl-3-(3,4-dichlorophenyl)-3-(3-[3-(homopiperazin-1-yl)azetidin-1-yl]propyl)piperidine;
and the pharmaceutically-acceptable salts thereof.

21. A pharmaceutical composition comprising a compound or salt according to any preceding claim together with pharmaceutically acceptable diluent, adjuvant or carrier.

22. A compound or salt according to any one of claims 1 to 20, or composition thereof according to claim 21, for use as a medicament.

23. The use of a compound or salt according to any one of claims 1 to 20 for the manufacture of a medicament for the treatment of a disease by producing an antagonist effect on a tachykinin acting at the human NK₁, NK₂ or NK₃ receptor, or a combination of two or more thereof.

24. The use of a compound or salt according to any one of claims 1 to 20 for the manufacture of a medicament for the treatment of an inflammatory disease such as arthritis, psoriasis, asthma or inflammatory bowel disease, a central nervous system (CNS) disorder such as anxiety, depression, dementia or psychosis, a gastro-intestinal (GI) disorder such as functional bowel disease, dyspepsia, irritable bowel syndrome, gastro-oesophageal reflux, faecal


incontinence, colitis, ulcerative colitis or Crohn's disease, a disease caused by Helicobacter pylori or other urease positive Gram negative bacteria, a urogenital tract disorder such as incontinence, hyperreflexia, impotence or cystitis, a pulmonary disorder such as chronic obstructive airways disease, an allergy such as eczema, contact dermatitis, atopic dermatitis, urticaria, eczematoid dermatitis or rhinitis, a hypersensitivity disorder such as to poison ivy, a vasospastic disease such as angina or Reynaud's disease, a proliferative disorder such as cancer or a disorder involving fibroblast proliferation, a fibrosing or collagen disease such as scleroderma or eosinophilic fasciitis, reflux sympathetic dystrophy such as shoulder/hand syndrome, an addiction disorder such as alcoholism, a stress-related somatic disorder, a peripheral neuropathy such as diabetic neuropathy, neuralgia, causalgia, painful neuropathy, a burn, herpetic neuralgia or post-herpetic neuralgia, a neuropathological disorder such as Parkinson's disease, Alzheimer's disease or multiple sclerosis, a disorder related to immune enhancement or suppression such as systemic lupus erythematosus, a rheumatic disease such as fibrositis, emesis, cough, acute or chronic pain, migraine, an ophthalmic disease such as proliferative retinopathy, ocular inflammation, conjunctivitis, a bladder disorder, or a viral disease such as influenza or a cold.

25. A method of treatment of a human or non-human animal which comprises producing an antagonist effect on a tachykinin acting at the human NK₁, NK₂ or NK₃ receptor, or a combination of two or more thereof, which comprises treating said human with an effective amount of a compound or salt according to any one of claims 1 to 20 or a composition according to claim 21.


26. A method of treatment of a human or non-human animal which comprises producing an antagonist effect on a tachykinin acting at the human NK₁, NK₂ or NK₃ receptor, or a combination of two or more thereof, which comprises treating said human with an effective amount of a compound or salt according to any one of claims 1 to 20 or a composition according to claim 21, where the disease is an inflammatory disease such as arthritis, psoriasis, asthma or inflammatory bowel disease, a central nervous system (CNS) disorder such as anxiety, depression, dementia or psychosis, a gastro-intestinal (GI) disorder such as functional bowel disease, dyspepsia, irritable bowel syndrome, gastro-oesophageal reflux, faecal incontinence, colitis, ulcerative colitis or Crohn's disease, a disease caused by Helicobacter pylori or other urease positive Gram negative bacteria, a urogenital tract disorder

such as incontinence, hyperreflexia, impotence or cystitis, a pulmonary disorder such as chronic obstructive airways disease, an allergy such as eczema, contact dermatitis, atopic dermatitis, urticaria, eczematoid dermatitis or rhinitis, a hypersensitivity disorder such as to poison ivy, a vasospastic disease such as angina or Reynaud's disease, a proliferative disorder such as cancer or a disorder involving fibroblast proliferation, a fibrosing or collagen disease such as scleroderma or eosinophilic fasciitis, reflux sympathetic dystrophy such as shoulder/hand syndrome, an addiction disorder such as alcoholism, a stress-related somatic disorder, a peripheral neuropathy such as diabetic neuropathy, neuralgia, causalgia, painful neuropathy, a burn, herpetic neuralgia or post-herpetic neuralgia, a neuropathological disorder such as Parkinson's disease, Alzheimer's disease or multiple sclerosis, a disorder related to immune enhancement or suppression such as systemic lupus erythematosus, a rheumatic disease such as fibrositis, emesis, cough, acute or chronic pain, migraine, an ophthalmic disease such as proliferative retinopathy, ocular inflammation, conjunctivitis, a bladder disorder, or a viral disease such as influenza or a cold.

27. A process for making a compound of formula (I) or salt thereof as defined in claim 1 comprising reaction of a compound of formula (II):




(II)



(III)

with a compound of formula (III), where X^1 , A, Ar^1 , X and R are as defined in Claim 1, and Z is a suitable leaving group.


28. A process for making a compound of formula (I) or salt thereof as defined in claim 1 comprising reductive amination of a compound of formula (IV) below,



(IV)

wherein X^1 , A and Ar^1 are as defined in claim 1, with an amine of the formula (III) as defined in claim 27, or a salt thereof.

29. A process for making a compound of formula (I) or salt thereof as defined in claim 1 comprising reaction of a compound of formula (VI) with a compound of formula (VII),




(VII)

(VI)

where X^1 , A, Ar^1 , X and R are as defined in Claim 1 and Z^1 is a suitable leaving group.


30. A process for making a compound of formula (I) or salt thereof as defined in claim 1 comprising reduction of a compound of formula (VIII):



(VIII)

where X^1 , A, Ar^1 , X and R are as defined in Claim 1.


31. A process for making a compound of formula (I) or salt thereof as defined in claim 1 comprising reaction of a compound of formula (IX),



(IX)

with a nucleophilic reagent acting as a RX^- "synthon" where X^1 , A, Ar^1 , X and R are as defined in Claim 1 and where Z^2 is a leaving group.


32. A process for making a compound of formula (I) or salt thereof as defined in claim 1 comprising reaction of a compound of formula (X):




(X)

with a reagent of formula X^1-A-NH_2 , where Z^3 and Z^4 are leaving groups and where X^1 , A, Ar^1 , X and R are as defined in Claim 1.

33. A process for making a compound of formula (I) or salt thereof as defined in claim 1 where R is




or




where Y is $NCO(C_{1-6}$ alkyl optionally substituted by one or more halogen), $NCO_2(C_{1-6}$ alkyl optionally substituted by one or more halogen), $NSO_2(C_{1-6}$ alkyl optionally substituted by one or more halogen), $NCOaryl$, NCO_2aryl , NSO_2aryl , $NSO_2(C_{1-6}$ alkyl optionally substituted by one or more halogen), NCH_2aryl , $N(C_{1-6}$ alkyl optionally substituted by one or more halogen), or $NCH_2(C_{3-7}$ cycloalkyl),

comprising reaction of a compound of formula (I) as defined in claim 1 where R is




or




as appropriate, and Y is NH, with a reagent of formula [(C₁₋₆ alkyl optionally substituted by one or more halogen)CO-, (C₁₋₆ alkyl optionally substituted by one or more halogen)CO₂-, (C₁₋₆ alkyl optionally substituted by one or more halogen)SO₂-, arylCO-, arylCO₂-, arylSO₂-, (C₁₋₆ alkyl optionally substituted by one or more halogen)SO₂-, arylCH₂-, (C₁₋₆ alkyl optionally substituted by one or more halogen)-, or (C₃₋₇ cycloalkyl)CH₂-]-Z⁵, where Z⁵ is a suitable leaving group.

34. A process for making a compound of formula (I) or salt thereof as defined in Claim 1 where R is




or



where Y is NH by reaction of a corresponding compound where Y is NCO₂C(CH₃)₃, with an acid.


35. A salt of the formula (V),



(V)


where X¹, A, Ar¹, X and R are as defined in Claim 1, and Y⁻ is the counterion of an acid HY.

36. A compound of formula (VI)




where Ar¹, X and R are as defined in Claim 1, or a salt thereof.

37. A compound of formula (VIII):



where X¹, A, Ar¹, X and R are as defined in Claim 1, or a salt thereof.

38. A compound of formula (IX),



where X¹, A, and Ar¹ are as defined in Claim 1 and where Z² is a leaving group, or a salt thereof.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/04177

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D401/06 C07D401/14 C07D211/22 C07D205/04 C07D403/04
A61K31/445

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 96 05193 A (PFIZER LIMITED) 22 February 1996 cited in the application see claims ---	1-38
A	EP 0 723 959 A (SANOFI) 31 July 1996 see claims ---	1-38
A	EP 0 512 901 A (ELF SANOFI) 11 November 1992 see claims ---	1-38
P, A	WO 97 25322 A (PFIZER RESEARCH AND DEVELOPMENT COMPANY N.V./S.A.) 17 July 1997 cited in the application see claims ---	1-38
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance
"E" earlier document but published on or after the international filing date
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
"O" document referring to an oral disclosure, use, exhibition or other means
"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
"&" document member of the same patent family

Date of the actual completion of the international search

1 October 1998

Date of mailing of the international search report

26/10/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Chouly, J

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/04177

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,A	<p>EP 0 791 592 A (PFIZER RESEARCH AND DEVELOPMENT COMPANY, N.V./S.A.) 27 August 1997 see claims</p> <p>-----</p>	1-38

INTERNATIONAL SEARCH REPORT

...ernational application No.

PCT/EP 98/ 04177

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 25, 26
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 25, 26
are directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 98/04177

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9605193	A	22-02-1996	AU 689303 B	26-03-1998
			AU 3254995 A	07-03-1996
			BR 9503582 A	30-04-1996
			CA 2197086 A	22-02-1996
			CZ 9700381 A	14-01-1998
			EP 0775132 A	28-05-1997
			FI 970523 A	07-02-1997
			HU 77771 A	28-08-1998
			JP 9508646 T	02-09-1997
			PL 318534 A	23-06-1997
			NO 970566 A	07-02-1997
EP 723959	A	31-07-1996	FR 2729951 A	02-08-1996
			JP 8245622 A	24-09-1996
			US 5656639 A	12-08-1997
EP 512901	A	11-11-1992	FR 2676055 A	06-11-1992
			AU 652046 B	11-08-1994
			AU 1591692 A	05-11-1992
			CA 2067877 A	04-11-1992
			CS 9201329 A	18-11-1992
			FI 921951 A	04-11-1992
			FI 951242 A	16-03-1995
			FI 951243 A	16-03-1995
			IL 101760 A	18-02-1997
			IL 117921 A	18-02-1997
			JP 5186425 A	27-07-1993
			MX 9202027 A	01-01-1993
			NO 178573 B	15-01-1996
			NZ 242586 A	26-10-1995
			RU 2083574 C	10-07-1997
			US 5770735 A	23-06-1998
			US 5625060 A	29-04-1997
			US 5340822 A	23-08-1994
WO 9725322	A	17-07-1997	AU 1195097 A	01-08-1997
			NO 982651 A	09-06-1998
EP 791592	A	27-08-1997	CA 2195923 A	28-07-1997
			JP 9208579 A	12-08-1997

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 98/04177

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 791592 A		US 5741790 A	21-04-1998